

PLANTARVM
VERTICILLATARVM
VNILABIATARVM
GENERALIA ET SPECIES

AUCTORE

IO. CHRISTIANO DANIELE SCHREBERO

SER. MARGGR. BNANDENS. ONOLD. ET CULMB. CONSIL. AVL
PHIL. ET MED. D. EIVSD. BOTAN. ET OECON. PROF. ORDIN. IN AGAM
ERLANG. ACAD. IMP. NAT. CVR. SOC. BOTAN. ET GEORGOPHIL
FLORENT. OECON. LIPSIENS. PATAVIN
VDINENS. SODAL



Dr. Steyermark

CVM ICONE AENEA

LIPSIAE

APVD SIGFR. LEB. CRYSIVM C¹CCCLXXVII



R. 11.391

β. Chamæpitys vulgaris folio trifido, floré roseo, lusitanica. TOVRN. inf. p. 208. exkluso synonymo, specimen 18 referendo.

H. in agris glareofis lapidofisque; in Franconia Brandenburgica circa Zwerniz prope Baruthum, in Saxonia passim circa Dresdam Lipsiamque; et in reliqua Germania, totaque Europa australiore inde a gradu $51\frac{1}{2}$, Palæstina, Africa boreali, Virginia; annua. Fl. post solstitium cum Hyssopo.

Plantula pulchra, vix dodratalis, pilosa. Fol. ima lanceolata, integra, trinervia, obtusa; paullo superiora lanceolata, tridentata; reliqua tripartita, avicularum pedes referentia, integrerrima, obtusa. Petioli infimi prælongi, superiores sensim decrescunt et dilatantur, trinervii. Flores parvi, brevissime pendunculati. Cal. depresso, ore contractior: dente summo duplo minore. Cor. lutea, labio serie dupli punctorum sanguineorum diviso. Filamenta basi barbata. — Varietas minor et tenerior est.

5. A. chia foliis trifidis, floribus axillaribus solitariis folio longioribus, caule diffuso.

Chamæpitys chia, folio trifido, magno flore. TOVRN & F. coroll. p. 34.

H. in insula Chio.

Simillima priori est et pariter pilosa, sed tenerior strictiorque. Flos multoties major, folium sibi subjectum longitudine excedit. Cor. lutea, labii crena profundiore.

6. A. Iva foliis linearibus antice dentatis, floribus axillaribus solitariis.

Teucrium Iva. LINN. sp. pl. 787. n. 7.

Chamæpitys moschata, foliis ferratis. BAVH. pin. 249. TOVRN. inf. p. 108. RAI. hist. p. 574. descr. GEOFFR. mat. med. 3. p. 307. descr. GARS. fc. 218.

Anthyllis altera. CLVS. hisp. p. 481. hisp. 2. p. 186. ic. bon.

Anthyllis chamæpityides minor. LOBEL. hist. p. 208. ic. 384. bon.

H. in arenosis, fuccis, lapidosis Galliae australis, Hispaniae, Lusitaniae CLVS. LOEFL. Siciliæ ad promontorium pachynum RAI. Corsicæ VALL. Mauritaniae SHAW. non igitur supra gr. 43. lat. Annua. Fl. post solstitium aestivum.

Herba pilis densis canis hispida, humistrata. Folia obtusiuscula, apicem versus paullo latiora cum dentibus duobus tribusve obtusis. Flores