



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789
<https://www.biodiversitylibrary.org/bibliography/4504>

v.3: <https://www.biodiversitylibrary.org/item/23434>

Page(s): Page 384, Page 385, Page 386

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 18 March 2024 7:16 AM
<https://www.biodiversitylibrary.org/pdf4/1682773i00023434.pdf>

This page intentionally left blank.

Cult. 1597. *Ger. herb.* 762. *f.* 1.

Fl. April—September.

H. ☉.

flexuosus.

9. *C.* foliis angulato-sublobatis, pomis cylindricis fulcatis curvatis. *Sp. pl.* 1437.

Serpent Cucumber, or Melon.

Nat.

Cult. 1597. *Ger. herb.* 763. *fig.* 3.

Fl. Most part of the Summer.

S. ☉.

B R Y O N I A. *Gen. pl.* 1093.

MASC. *Cal.* 5-dentatus. *Cor.* 5-partita. *Filamenta* 3.

FEM. *Cal.* dentatus. *Cor.* 5-partita. *Stylus* 3-fidus.

Bacca subglobosa, polysperma.

latebrosa.

1. *B.* foliis subtrilobis pilosis basi attenuatis.

Hairy Briony.

Nat. of the Canary Islands. Mr. Francis Masson.

Introd. 1779.

Fl. June.

G. H. 4.

OBS. Facillime distinguitur a congeneribus, foliis basi minime cordatis, sed secundum petiolos subdecurrentibus.

alba.

2. *B.* foliis palmatis utrinque calloso-scabris. *Sp. pl.* 1438.

White Briony.

Nat. of Britain.

Fl. May.

H. 4.

palmata.

3. *B.* foliis palmatis lævibus quinquepartitis: laciniis lanceolatis repando-ferratis. *Sp. pl.* 1438.

Palmated Briony.

Nat. of Ceylon.

Introd.

Introd. 1778, by Messrs. Gordon and Græfer.

Fl. S. 4.

4. *B. foliis cordatis angulatis glabriusculis basi subtus glandulosis, cirrhis simplicibus.* *grandis.*

Bryonia grandis. *Linn. mant.* 126.

Great-flower'd Briony.

Nat. of India.

Introd. 1783, by Mr. John Græfer.

Fl. S. 4.

5. *B. foliis cordatis angulatis lobatisque calloso-hispidis, cirrhis simplicibus, baccis globosis, seminibus muricatis.* *scabrella.*

Bryonia scabrella. *Linn. suppl.* 424.

Bristly Briony.

Nat. of the East Indies. *John Gerard Koenig, M. D.*

Introd. 1781, by Sir Joseph Banks, Bart.

Fl. May—July. S. 6.

6. *B. foliis cordatis angulatis subtus villosis; supra calloso-scabris, cirrhis simplicibus, baccis globosis, seminibus lævibus.* *scabra.*

Bryonia scabra. *Linn. suppl.* 423.

Globe-fruited Briony.

Nat. of the Cape of Good Hope. *Mr. Fr. Masson.*

Introd. 1774.

Fl. September and October. G. H. 4.

7. *B. foliis cordatis angulatis supra venisque subtus calloso-scabris: callis remotis, cirrhis simplicibus, baccis globosis.* *verrucosa.*

Rough Briony.

Nat. of the Canary Islands. *Mr. Francis Masson.*

Introd. 1779.

Fl. G. H. 4.

- africana.* 8. *B. foliis palmatis quinquepartitis utrinque lævibus: laciniis pinnatifidis. Sp. pl. 1438.*
 African, or Smooth-leav'd Briony.
Nat. of the Cape of Good Hope.
Cult. 1759, by Mr. Ph. Miller. *Mill. dict. edit. 7. n. 2.*
Fl. July and August. G. H. 4.

S I C Y O S. *Gen. pl. 1094.*

MASC. *Cal.* 5-dentatus. *Cor.* 5-partita. *Filamenta* 3.
 FEM. *Cal.* 5-dentatus. *Cor.* 5-partita. *Stylus* 3-fidus.
Drupa 1-sperma.

- angulata.* 1. *S. foliis angulatis. Sp. pl. 1439.*
 Angular-leav'd Sicyos, or Single-seeded Cucumber.
Nat. of North America.
Cult. 1732, by James Sherard, M. D. *Dill. elth. 58.*
t. 51. f. 59.
Fl. July—September. H. ☉.

G Y N A N D R I A.

A N D R A C H N E. *Gen. pl. 1095.*

MASC. *Cal.* 5-phyllus. *Cor.* 5-petala. *Stam.* 5, *Styli*
 rudimento inserta.
 FEM. *Cal.* 5-phyllus. *Cor.* 0. *Styli* 3. *Caps.* 3-locularis.
Sem. 2.

- Telephioides.* 1. *A. procumbens herbacea. Sp. pl. 1439.*
 Bastard Orpine.
Nat. of Italy and the Levant.
Cult. 1732, by James Sherard, M. D. *Dill. elth. 377.*
t. 282. f. 364.
Fl. July and August. H. ☉.