



<https://www.biodiversitylibrary.org/>

**Linnaea : Ein Journal für die Botanik in ihrem ganzen
Umfange**

Berlin F. Dümmler 1826-1882

<https://www.biodiversitylibrary.org/bibliography/626>

Bd.38=NF Bd.4 (1874):

<https://www.biodiversitylibrary.org/item/10883>

Page(s): Page 699, Page 700

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 19 March 2024 7:34 AM

<https://www.biodiversitylibrary.org/pdf4/1683154i00010883.pdf>

This page intentionally left blank.

Notulae in Campanulaceas herbarii regii berolinensis.

Auctore

W. Vatke.

1. *Jasione corymbosa* C. Koch. Beitr. Fl. or. in Linnaea XIX p. 30 est *J. supina* Sieber. Cl. auctor suam infauste cum *J. montanae* var. *arenaria* (= var. *littoralis* Boiss. voy., quae nunc *J. blepharodon* Boiss. et Reut.) et cum *J. corymbosa* Poir., specie altera hispanica, altera boreali-africana satis distinctis junxit.

2. *Lightfootia sessiliflora* (L.) Spr. nomen nuper a cl. Sonder restitutum jam a Sprengelio datum (syst. veg. I p. 809), qui quidem postea in herbario zeyheriano stirpem diversam cum hac commiscuit.

3. *L. thunbergiana* Buek. In montibus Hantam Africae australis legit cl. Dr. Meyer 1869.

4. *Cephalostigma erectum* (Rth.) Vatke. Dentella e. Rth., *D. perotifolia* Willd. herb., *Wahlenbergia* p. W. et A., *C. hirsutam* Edgew., *C. Schimperii* Hochst. Cf. Hooker f. et Th. in Journ. Linn. Soc. II. 9.

Synonymis addendum adhuc: *Wahlenbergia Schimperii* Hochst. jam ab ipso cl. auctore rite huc relatum, sed a Schweinfurthio lapsu iterum sejunctum.

5. **Treichelia longibractea** (Buck) **Vatke**. *Leptocodon* (non Hook. f. et Th. Journ. Linn. Soc. II. p. 17, 1858) l. Sonder Fl. cap. III. p. 584 (1864—5) Genus dictum cl. A. Treichel, unionis botanicae berlinensis librorum praefecto mihiq̄ amicissimo.

6. **Canarina canariensis** (L) **Vatke**. (*C. Campanula* Lam) Nomen linnaeanum restituendum.

7. **Wahlenbergia pilosa** **Buck**. In montibus Hantam coll. Dr. Meyer. Exemplaria pygmaea, ceterum cum descriptione exacte convenit.

8. **W. oxyphylla** **DC. f.** Idem legit ibidem.

9. **W. riparia** (Lepr. et Perr.) **DC. f.** huc Barter in Baikie's Niger Expedition 1857—9 n. 1202 Cum spec. orig. h. reg. berol. omnino congruit.

10. **W. Mannii** **Vatke**. *W. polyclada* Hook. f. Journ. Linn. soc. VI. p. 15 (non **CD. f.**).

11. **W. nudicaulis** **DC. f.** Exempl. e mont. Hantam cl. Meyer cum descriptione candolleana praeter folia pilosiuscula dicta (in nostra hispida) plane conveniunt.

12. **W. Krebsii** **Cham. in Linnaea VIII** (1833) nomen *W. Zeyheri* **Buck** in E. et Z. en. antiquius restituendum.

13. **W. campanuloides** (Del.) **Vatke**. (*W. cervicina* D. Cf.).

14. **W. etbaica** (**Schweinf.**) **Vatke**. (*Laurentia* e. Schweinfurth in herb. reg. berlin.) annua caule subsimplici, ad medium folioso, superne nudo, inferne hispidissimo superne glabrato, foliis alternis ovato-lanceolatis basi attenuata sessilibus, superioribus acutis, subtus nervo medio hispidis, floribus parvis terminalibus lateralibusque remotis, tubo calycis glabri elongato gracili, laciniis tubum vix aequantibus lanceolatis acutis integris. ☉

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 699]

Ka EEE

Notulae in Campanulaceas herbarii regüi
berolinensis,

Auctore

W. Vatke.

1. Janlone corymbosa C. Koch, Beitr. Fl. or. in
Linnaea XIX p. 30 est J. supina Sieber. Cl, auctor suam
infauste eum), montanae var. arenaria (= var, littoralis
Boiss. voy., quae nunc J. blepharo don Boiss. et Reut.) et
eum J. corymbosa Poir., specie altera hispanica, altera
boreali-africana satis distinctis junxit.

2%. Lightfootia sessilillora (L.) Spr. nomen nuper a
el. Sonder restitutum jam a Sprengelio datum (syst. veg. I
p- 809), qui quidem postea in herbario zeyheriano stirpem
diversam cum hac commisit,

3. L. thunbergiana Buck. In montibus Hantam A-
friae australis legit cl. Dr. Meyer 1869.

4. Cephalostigma erectum (Rth.) Vatke, Dentelia

" e& Rth., D. perotifolia Willd. herb., Wahlenbergia p. W. ei
&., C. hirsutam Edgew., Q. Schimperi Hochst. Cf. Hooker
f. et Th. in Journ. Linn. Soc. il, 9.

Synonymis addendum adhuc: Wahlenbergia Schimperi
Hochst. jam ab ipso el. auctore rite huc relatam, sed a
Schweinfurthio lapsu iterum sejunctam.

[Begin Page: Page 700]

700

w . 5. Treichelia longibracteata (Buck) Vatke. Lep-
"0° teeoden (nom Hook. f, et Th. Journ. Linn. Soc. I. p. 17,
1858) 1. Sonder Fl. cap. II. p. 584 (1864—5) Genus dia-
tum cl. A. Treichel, unionis botanicae berolinensis lihorum
praefecto mihique amieissimo. j

6. Canarina canariensis (L) Vatke. (C. Campanula
Lam) Nomen linnaeanum restitendum. "

9. Wahlenbergia pilosa Buck. In montibus Hantam
soll. Dr. Meyer. Exemplaria pygmaca, ceterum cum descrip-
, tione exaete convenit.

8 *W. oxyphylla* DC. f. Idem legit ibidem.

9. *W. riparia* (Lepr. et Perr.) DC. f, huc Barter in Baikie's Niger Expedition 1857—9 n. 120% Cum spec. orig. hu reg. berol. omnino congruit,

10. *W. Mannii* Vatke. *W. polyclada* Hook. f. Journ. Linn. soe. VL p. 15 (non CD. f.).

il. *W. nudicaulis* DC. f, Exempl. e mont, Hautam el, Meyer cum descriptione candolleana praeter folia pilosiuscula dieta (in nostra hispida) plane conveniunt.

12. *W. Krebsii* Cham, in Liunsea VEIL (1833) nomen *W. Zeyheri* Baek in E, et Z..en. antiquins restituendum. . "

13. *W. campanuleides* (Del.) Vatke. {*W. cerviciua* D.Ck.).

14. *W. etbaica* (Schweinf.) Vatke. (Laurentia e, Schweinfurth in herb. reg. berlin.) aunua caule subsimplici, ad medium folioso, superne nado, inferne hispidissime superne glabrato, foliis alteruis ovato-lanceolatis basi attenuata sessilibus, superioribus acutis, subtus nervo medio hispidis, Horibus parvis terminalibus lateralibusque remotis, tubo calyis glabri elongato gracili, laciniis tubum vix aequantibus | lauceolatis acutis integris. @