



<https://www.biodiversitylibrary.org/>

**Caroli Linnaei ... Species plantarum :exhibentes plantas  
rite cognitatas, ad genera relatas, cum differentiis specificis,  
nominibus trivialibus, synonymis selectis, locis natalibus,  
secundum systema sexuale digestas**

Holmiae, Impensis Direct. Laurentii Salvii, 1762-63

<https://www.biodiversitylibrary.org/bibliography/11179>

**v.2:** <https://www.biodiversitylibrary.org/item/43462>

Page(s): Text, Title Page, Page 1151

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 14 July 2024 5:12 AM

<https://www.biodiversitylibrary.org/pdf4/1716087i00043462.pdf>

This page intentionally left blank.

CAROLI LINNÆI

*Equitis aur. de Stella polari,*

ARCHIATRI REGII, MED. & BOTAN. PROFESS. UPSAL. ACAD.  
PARIS. UPSAL. HOLMENS. PETROPOL BEROL. IMPERIAL.  
BERNENS. LONDIN. ANGLIC. EDENBURG. MONSPEL.  
TOLOS, FLORENT. SOC.

SPECIES  
PLANTARUM,

EXHIBENTES  
PLANTAS RITE COGNITAS,  
AD

GENERA RELATAS,  
CUM

DIFFERENTIIS SPECIFICIS,  
NOMINIBUS TRIVIALIBUS,  
SYNONYMIS SELECTIS,  
LOCIS NATALIBUS,  
SECUNDUM

SYSTEMA SEXUALE  
DIGESTAS.

MISSOURI  
BOTANICAL  
GARDEN.

TOMUS II.

Editio Secunda, aucta.

Cum Privilegio S. R. M:tis Sveciæ & S. R. M:tis Poloniæ ac Electoris Saxon.

HOLMIÆ  
IMPENSIS DIRECT. LAURENTII SALVII

1763.

Linnaeus  
QK91  
- C500  
1762-63  
v. 2

A.

6. **CARDUUS** foliis decurrentibus dentatis: margine palustris, spinosis, floribus racemosis erectis, pedunculis inermibus.

*Carduus* foliis linearibus dentatis spinosissimis, caule crispo spinoso, floribus erectis. *Roy. lugab.* 134. *Fl. svec.* 659; 720. *Dalib. paris.* 249.

*Carduus palustris* *Baub. pin.* 377. *prodr.* 156.

*Carduus spinosissimus erectus angustifolius palustris.* *Moris. hist.* 3. p. 153. f. 7. t. 32. f. 13.

*Carduus polycephalus angustifolius.* *Dalech. hist.* 1743.

*Habitat in Europæ pratis subpaludosis.* 4.

*Calyx arcte imbricatus, laevis, mucronibus vix conspicuis.*

7. **CARDUUS** foliis decurrentibus pinnatifido-sinuatis pycnocephalus, pubescentibus spinosis, pedunculis nudis tomentosis, calycibus deciduis.

*Carduus nemorosus italicus.* *Barr. obs.* 925. t. 417.

*Carduus pycnocephalus sylvestris.* *Triumph. obs.* 100. t. 101.

*Habitat in Europa australi.* 4.

*Planta quasi media inter C. crispum & Onopordon.*

*Caulis pedalis s. altior, albido-pilosus, crispus, totaliter foliorum marginibus utriusque decurrentibus, alternatim flexuosis spinosisque. Folia Radicalia sublanceolata, subsinuata; Caulina oblonga, semipinnatifida, dentata, utrinque pubescentia, subtus albidiora, tomentosa s. sublanata. Pedunculi breves, tomentoso-albi.*

*Calyces magnitudine avellanæ, oblongi, imbricati squamis subulatis, erectis, patulis, subspinosis. Corolla purpurea limbo erecto, magnitudine Serratulæ tinctoriæ. Differt a C. acanthoide pedunculis nudis nec alatis. Calyces maturi toti decidunt.*

8. **CARDUUS** foliis decurrentibus lanceolatis, denticulis dissectis, inermibus, calyce spinoso. *Hort. cliff.* 392. \*

*Dalib. paris.* 249.

*Cirsium anglicum.* *Lob. ic.* 583. *Dalech. hist.* 584. *figura tenuis.*

*Cirsium majus, singulari capitulo magno, s. incanum varie dissectum.* *Baub. pin.* 377.

*Habitat in Anglia, Gallia.* 4.

*Folia carent omnino aculeis, diversus itaque a C. heterophyllo & betenioides.*

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Text, Title Page]

f

1

CAROLI LINKJBI

. iTti

Equitis aur. de Stella folari<sup>^</sup> <sup>^</sup> 2L

ARCHIATRI ReGH, MeD & BoTAN. PROFESS. IJPSAL. AcAD.

Faris. Uj>sal. Holmens. Petropol Berol Imperial.

B&RNENS. LONDIN AnGLIC. EdENBURG. MoNSPEL,

ToLos, Fi; ORENT. Sbc.

SPECIES

M

\$

EXHIBENTES

PLANTAS RITE COGNITAS

AD

GENERA RELATAS,

CUM .

DIFFERENTIIS SPECIFICISj

NO MINIBUS TRIVIALIBUS,

SYNONYMIS SELECTIS,

LOCIS NATALIBUS,

SECUNDUM

ST ST EM A SE XVAL B

\_,^, DIGESTAS.

>

BOTANICAL,

^ \RDEN. «

TOMUS

f

Editio Scv»qd45 aufta.

Cum Prhikgio S. R M'.tis, Svfcia^ \$, jR, M:tis PtloJii^ ac EhSofit Saxoi^.

HOLMIM

Impensis Direct, LAURENTII SALVII

1763.

**[Begin Page: Page 1151]**

f

SYNGENESIAiPOLYGAMIA^QUALIS. 1151

6. CARDUUS foliis decurrentibus dcntatis : marginepilulfrls.

fpinafis, floribus racemoiis cre6lis, pcdunculis i- .

\*^ncrmibus.

Carduus foliis Hnearibus c^entatls fpinofiffimis-. caule

cripfo fpinofo, tiolibus erc6lis. RoyJugab. 134. tL

Jvec, 659; 720. Dalti;. pari/. 249.





Caulis pedalis J. alitor J albid<sup>^</sup>-pilc<sup>fiis</sup><sup>^</sup> Ei ifpus, t<sup>^</sup>aliter  
jstior<sup>km</sup> riiarginibus utriitque decurremib<sup>Hs</sup> <sup>^</sup> alter<sup>^</sup>  
natimfiexuofts fpinofuque. <sup>^</sup>oWjkKadicaltafublanceo<sup>^</sup>  
lata-<sup>^</sup>fubfinuata ; Cuiilina oblanga<sup>^</sup>fimipinnatijida<sup>^</sup> den<sup>^</sup>  
fata;<sup>^</sup> tutri<sup>^</sup>jque pubtfcentia<sup>^</sup> fubtus albidiora<sup>^</sup> t<sup>^</sup>men<sup>^</sup>  
tofa f. fublanata. Pcdunculi breves<sup>^</sup> tomentofj-albi.  
Calyces ntiignitudiMe avellana <sup>^</sup> oblongi <sup>^</sup> imbricati

\

fquamis fubu<sup>Utis</sup> <sup>^</sup> erects \ patu<sup>Us</sup> <sup>^</sup> fubjftnofs. Qo<sup>^</sup>  
toW'X pnrpurea limbo ereHo <sup>^</sup> tfiagnitudine Serratula  
tinSorie. Dirfert a C. acanthoide pedmneu<sup>Us</sup> mudis

vi€ a.latts. Calyces maturi toH 4ecidupst.

y

8, CARDtJtJS foliis decurrentlbas iancoiatls<sup>^</sup> dcaticu<sup>^</sup>ifTeftus-  
Us inermibus, calyce fpinoib. K(>r/. atff-, 3<sup>^</sup>-, \*<sup>^</sup>

Dalib. parif, 249;

Cir{ium anglicuin. L<sup>^</sup>b.ic. 5S3. Daleck hifl, <sup>^</sup>4. fi\*

gura tenus,

Cirfium niajuS, fingulari capitalo magno, f. incanum

varie difTcdum. Bauh. pin, 577. ,

Hakitat in Anglia, Gallia. %.

Fulia (arent (^mnino acuUis^ diverfus itaque a Ci he-^

terophyllQ ^ hehmQidi.

\

9. CAR-

N.

1