



<https://www.biodiversitylibrary.org/>

**Synopsis plantarum succulentarum :cum descriptionibus,
synonymis, locis, observationibus anglicanis, culturaque**

Londini, Typis R. Taylor et Socii, 1812

<https://www.biodiversitylibrary.org/bibliography/9462>

Item: <https://www.biodiversitylibrary.org/item/38208>

Page(s): Page 13, Page 14

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 20 March 2024 11:00 AM

<https://www.biodiversitylibrary.org/pdf4/1683571i00038208.pdf>

This page intentionally left blank.

HABITAT C. B. S.

S. h.

FLORET Jul.—Nov.

INTROD. 1795. Hort. kew. l. c.

DIGYNIA.

SARCOSTEMMA.

Asclepiadea. Massæ Pollinis læves, 10, pendulæ. Corona staminea duplex: exterior cyathiformis vel annularis, crenata. Cor. rotata. Hort. kew. ed. 2. v. 3. p. 76.

S. (Twisting) H. Kew. l. c.—*Cynanchum viminale*, *caule volubili perenni aphylllo*. Pers. 1. syn. 272.—*Euphorbia viminalis*. Linn. sp. + pl. 2. 649.—*Tithymalus* g. Mill. dic. ed. 1.

HABITAT in Africæ maritimis et Indiæ nemorosis. Caulis nudus, crassitudine calami. Pers. l. c.

S. h.

FLORET

CULT. 1731.

CEROPEGIA.

Contorta. *Folliculi* 2, erecti. Sem. plumosa.

Cor. urceolatæ limbus connivens. Pers. syn. 277.

C. (Forked) tota carnosa; caulis strictis dichotomis articulatis teretibus, foliis linearibus ma- acutis horizontalibus sesquiuncialibus; supra 1. costa venisque simplicibus albis. *

HABITAT

S. h.

FLORET Aug.

INTROD. ante 1806.

C.? (Leafless) carnosa, ramis dichotomis mul- *aphylla*. tiarticulatis debilibus teretibus; foliolis stipu- 2. læformibus adpressis, mox emarcidis. *

Pentandria Dig. Ceropegia.

HABITAT C. B. S.? S. h.

FLORET

CULT. circa 1808.

OBS. *Foliola* circa lineam longa, et demum fere deltoidea, et nisi lente fere inconspicua. — Parum lactescit, habitu omnino *Tithymali* denudati, vel potius *Cacti* penduli *Linn.*

HOYA.

Asclepiadea. Massæ Pollinis læves, 10, erecto-conniventes. *Antheræ* membrana terminatæ. *Corona staminea* 5-phylla; foliolis depresso-angulo interiore producto in dentem antheræ incumbentem. *Cor. rotata.* *Hort. kew. ed. 2.* 84.

carnosa. H. (Fleshy) foliis ovali-oblongis carnosis, corollis barbatis, coronæ foliolis subtus sulcatis. *Hort. kew. l. c.—Asclepias carnosa.* *Bot. mag. 788.—Smith's sexot. bot. t. 70.*

HABITAT in Asia. S. h.

FLORET Jul. Aug.

INTROD. 1802.

OBS. It will survive, but not flower in the Greenhouse: and in a strong heat, and poor dry soil flowers best.

STAPELIA. Gen. pl. 432.

Ligulæ (s. *Nectarii stella exterior*) 5, basi connatae, imo corollæ insertæ, genitalia ambientes. *Stamina* 5, corniculiformia sæpissime bipartita (exteriore ramo incurvo-cultriforme, interiore recurvo-calcariforme) intra *ligulas*, et eas basi connata, interne ipsa cava basi, anthers duas distinctas sessiles cerinas (recon-

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 13]

Peniandria Monog. . Echites. ME
HanrrAT C. B. S. : Lore das
FronET Jul.—Nov.
IxNThOD. 1795. Hort. kew. I. c.
DIGYNI 4.
-. SARCOSTEMMA.
B6 rue JMassce Pollinis leves, 10, pen-
du Corona staminea duplex: exterior
eritiiionio n pague, crenata. . Cor.
rotata. Hort. kew. ed. 2. 8.2. 70.
S. (Twisting) H. Kew. L6. s Dynicli vimi- vimingle.
ale, caule volulili perenni aphyollo. - Pers. — 1.
syn. Sica viminalis. Linn.sp. . t
pl. 2. 649.—Tithymalus 9. Mill. dic. ed. 1.
HaBrTAT in Africee maritimis et Indize nemo-
rosis. — Caulis nudus, crassitudine calami.
Pers. I. c. oh.
FLoRET ...
CurT. ama.
CEROPEGIA.
don. Folliculi 9, erecti. Sem. WR
. Cor. urceolatze limbus counivens. Pers. syn.
277.
C. (Forked) tota carnosa ; caulibus strictis di- dickoto- z
chotomis articulatis teretibus, foliis limearibus ma.
acutis horizontalibus sesquiuncialibus; supra l.
costa venisque simplicibus albis. »
HántraT 7.2. 8. 5h.
FronET Aug.
GC - Fide) carnosa, ramis Icbowale mul- péglis
tis debilibus teretibus ; uh Pu s.
eter us adpressis, mox emareidis

[Begin Page: Page 14]

carnosa.
1l.
Pontisdria Digi: sidmp
—n B. S.? : S. bh.
FLoRET ...
Cin. circa 1808.
Oss. Foliola circa lineam longa, et 'demum
fere deltoidea, et misi lente fere inconspicua.
Parum lactescit, habitu omnino Tithymali
denudati, vel pitié Cacti viu Linn.
HOYA.
ou Masse Pollinis leves, 1o, erecto-
conniventes. Zntherc membrana terminatz.
' Corona staminea 5- pbi: foliolis Ate;

angulo i interiore producto in dentem antherze
e umbentem. Cor. rotata. Hort. kew. ei
. 84.

H. FANS foliis ovali-oblongi i\$, co-
rollis barbatis, coronz foliolis oicork. sulcatis.
Hort. kew. l. c.— Asclepias carnosa. ^ Bot.
mag. 788. SRL SOC 1 lot. t. 70.

HaBrrAT jn'Asla; .— Sy.

FLónET Jul. Ee

INTROD.!

Oss. It wil survive, but not dieup«s in the
Greenhouse : and in a strong heat, and poor
: A soil flowers best,
STAPELIA. Gen. pl. 439.

Ligule (s. Nectarii es exterior! 5, basi connat^a
tz; imo corollze i insertze, genitalia. ambientes.
amina 5, cornicuhformia. Ssepissime. bipar-
tita (exteriore ramo. incurvo-eultriforme e, Los
BU recurvo-calcarifi ligülc
i àsi connata, interne i i psa cava à
- theras duas distinctas sessile