



<https://www.biodiversitylibrary.org/>

**Supplementum plantarum Systematis vegetabilium  
editionis decimae tertiae, Generum plantarum editionis  
sextae, et Specierum plantarum editionis secunda**

Brunsvigae, Impensis Orphanotrophei, 1781

<https://www.biodiversitylibrary.org/bibliography/555>

**Item:** <https://www.biodiversitylibrary.org/item/10321>

Page(s): Page 135, Page 136

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 22 March 2024 2:54 AM

<https://www.biodiversitylibrary.org/pdf4/1684079i00010321.pdf>

This page intentionally left blank.

*Folia* majora, longius petiolata, minus acuminata,  
quam in *L. glabro*.

*Pedunculi* axillares & terminales, erecti, terminantes  
in umbellas.

*Umbellæ* simplices, paucifloræ, subinvolutæ: fo-  
liis binis s. ternis.

*Corolla* campanulato - infundibuliformis, lutea.

### Observatio.

Proprium hujus generis est Capsula bilocularis, bival-  
vis: valvularum marginibus intortis, & Stigma  
bilamellatum.

#### *repens.* SHEFFIELDIA.

*Sheffieldia repens.* Forst. gen. n. 9.

*Habitat* in Nova Zeelandia.

Parvula, procumbens & repens, Peplidi Portulæ adeo  
similis, ut demta fructificatione eandem cre-  
deres.

*subloba-* CONVOLVULUS procumbens, foliis superiori-  
*tus.* bus apice dentato - repandis, floribus capitatis.

*Habitat* in India. ♂.

*Folia* subpilosa.

*inferiora* cordata.

*superiora* obtusa, sublobato - repanda in apice.

*Flores* aggregati, terni, sessiles, paralleli.

*Involucrum* hexaphyllum.

*Corollæ* magnæ.

*scoparius.* CONVOLVULUS fruticosus, erectus, glaber,  
ramis virgatis, foliis sessilibus linearibus, racemo  
terminali, pedunculis subtrifloris.

Lena-Noel s. Lignum Rhodium. Barranc.

*Habitat* in Barrancas circa oppidum St. Crux.

Fr. Masson. ♂.

*Facies* Spartii s. Genistæ & non Convolvuli.

*Caulis* teres, glaberrimus.

*Rami* simplicissimi, errecti, foliosi.

*Folia* alterna, remota, brevia, gracilia, linearia, sub-  
pilosa, errecta.

*Racemus terminalis, subsecundus.*

*Pedunculi alterni, solitarii, remoti, plerumque triflori, rarius uniflori, bracteati.*

*Corollæ extus hirsutæ, albæ.*

*Lignum album, durum: striis radiantibus.*

*Rasura hujus ligni spargit odorem rosaceum & saporem acriuscum habet. An verum Lignum Rhodium Officinarum?*

*floridus.* CONVOLVULUS fruticosus, prostratus, canus, foliis linear - lanceolatis glabris, thyrso terminali pyramidali composito.

*Habitat in rupibus circa Barrancas. Franc. Mafson. ḫ.*

*Similis C. Cneoro.*

*Caulis lignosi, procumbentes.*

*Rami erecti, stricti, cani.*

*Folia petiolata, linearia, basi attenuata, integra, undulata, digitalia, venosa, glabra.*

*Pedunculi ex axillis foliorum superiorum, ramosi: ramis ramosis; sic continuando formatur panicula magna, ovata s. thyrsoidea.*

*Flores numerosissimi, minores.*

*Calycis foliola ovata, concava, acuta.*

*Corollæ dilute rubræ, extus hirsutæ.*

*Copia florum omnium pulcherrimus.*

*grandiflorus.* CONVOLVULUS foliis cordatis ovatis obtusiusculis integerrimis, pedunculis subbifloris, calycibus coriaceis, caule petiolisque pubescentibus.

Convolvulus maximus, semine hirsuto. Herm. mis.

Munda - Valli. Rheed. mal. v. II. p. 103. t. 50.

*Habitat intra spinetas ad littora fluminum Tanschur. König. ḫ.* H.U.

*Caulis arboreus, primo erexitus, dein volubilis, pubescens.*

*Folia magna, cordata: lobis rotundatis; obtusa, glabra, petiolata.*

*Petioli pubescentes.*

*Pedunculi axillares, plerumque biflori.*

*Flores majores.*

*Calyx di - l. triphyllus, coriaceus.*

CONVOL-

**The following text is generated from uncorrected OCR or manual transcriptions.**

**[Begin Page: Page 135]**

sho on  
i T e T T fog ow foliis fuperiori-  
; bus : apice denara.- gpi, floribus popan  
Habitat in India. .  
PEE IL  
A:  
ndi X  
DN a

**[Begin Page: Page 136]**

extus mn NES  
cili yrs egi: Ce M NV  
Šo us pis etiolata, x bafi at bl, integra, un-  
| Flores mayores