



<https://www.biodiversitylibrary.org/>

**Supplementum plantarum Systematis vegetabilium  
editionis decimae tertiae, Generum plantarum editionis  
sextae, et Specierum plantarum editionis secunda**

Brunsvigae, Impensis Orphanotrophei, 1781

<https://www.biodiversitylibrary.org/bibliography/555>

**Item:** <https://www.biodiversitylibrary.org/item/10321>

Page(s): Page 131

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 8 March 2024 12:55 PM

<https://www.biodiversitylibrary.org/pdf4/1678120i00010321.pdf>

This page intentionally left blank.

*candi-*  
*cans.* ECHIUM fruticosum, hirsutum, albicans, foliis lanceolatis utrinque acuminatis venoso - nervosis, panicula terminali, pedicellis spicatis.

*Habitat* in Maderæ rupibus altis. *F. Masson.* Ḥ.  
*Frutex* pulcher.

*Caules & Rami* lignosi, subtomentosi : Cicatricibus foliorum linearibus, transversalibus, approximatis.

*Folia* spithamæa, approximata, lanceolato - subulata, basi attenuata usque ad insertionem ubi dilatata, supra lineata, subtus venosa: venis longioribus simplicibus.

*Folia* paniculam versus imbricata, basi angustiora.

*Panicula* ramulum terminans, conica, multispicata.

*Spicæ* pedicellatæ, simplices, secundæ, incurvatae:

*Corollæ* ex minoribua hujus generis, subhirsutæ, cæruleæ.

*Tota planta* albicat, quasi argentea: hirsutie albida, approximata, molliori, omnes partes, vix corolla excepta, tegente.

*gigan-*  
*teum.* ECHIUM fruticosum, ramis canis glabris, foliis linear - lanceolatis scabriusculis sessilibus, thyrso terminali: spicis simplicissimis.

*Habitat* in rupibus Teneriffæ. *Franc. Masson.* Ḥ.  
Altissimum.

*Rami* lignosi, teretes, cani, glabri.

*Folia* linearia, spithamæa, latitudine digitæ, scabra, venosa.

*Thrysus* terminalis, maximus, pyramidalis, constans ex pedunculis longis, spicatis.

*Flores* secundi.

*Corollæ* obtusæ, subæquales, albæ.

*strictum.* ECHIUM fruticosum, foliis petiolatis ellipticis hispidis scabris, caule reverse - hispido, pedunculis axillaribus spicato - capitatis.

*Habitat* in rupibus Teneriffæ. *Masson.*

Distinguitur caule stricto, foliis petiolatis oblongis, pilis caulis deorsum versis.

*Pedunculi* axillares in summitate caulis solitarii, nudæ, apice spiculis tripartitis, coarctatis in capitulum.

*Flores* parvi, cærulei.

*Stamina* exserta.