



<https://www.biodiversitylibrary.org/>

**Prodromus systematis naturalis regni vegetabilis, sive,
Enumeratio contracta ordinum generum specierumque
plantarum huc usque cognitarium, juxta methodi naturalis,
normas digesta**

Parisi, Sumptibus Sociorum Treuttel et Wurtz, 1824-73

<https://www.biodiversitylibrary.org/bibliography/286>

v.15:pt.2 (1862-1866): <https://www.biodiversitylibrary.org/item/7166>

Page(s): Page 108

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 25 March 2024 12:02 PM

<https://www.biodiversitylibrary.org/pdf4/1684839i00007166.pdf>

This page intentionally left blank.

et 1510!). Vulgo *Tabayba dulce*. — Webb phyt. Can. t. 209 et tab. phytost. t. 6. In Senegambiæ regno Cayor ad sepes formandas culta sub nomine Selane (Heudelot!), an ibi vere spontanea? Frutex 4-5-pedalis. Folia 9-12 lineas longa, 2-2½ lata. Involucra quædam hermaphrodita, sæpius mascula. Capsula piso major. Succus mitis et jucundus. Species a sequentibus inflorescentiâ monocephalâ et defectu foliorum floralium distinctissima. (v. s.)

419. *E. MELLIFERA* (Ait. hort. Kew. ed. I, vol. 3, p. 403), glabra, arborescens, ramis superne foliosis, cymis pedunculatis paniculam terminalem simplicem vel compositam basi tantum foliosam formantibus, foliis basi attenuatis elongato-lanceolatis acutis mucronatis subsessilibus nervis secundariis manifestis, foliis floralibus ad basin involucrorum binis oblongis concavis mucronatis ciliatis, involucri campanulati lobis late et breviter ovatis ciliatis, glandulis quinque semiorbiculatis subtus hirtulis, stylis longis ad medium coalitis bifidis apice capitatis, capsulæ magnæ sphericæ subtrilocæ sparsim tuberculatæ coccis dorso rotundatis, semine ovato lævi dorso carinato, carunculâ patellari semilunatâ. 5 In sylvaticis Maderæ (Webb! Mason!). — Vent. Malm. t. 30, Bot. mag. tab. 1305. *E. longifolia* Lam. dict. II, p. 417. Arbuscula 6-15-pedalis. Folia 6-12 poll. longa, 10-15 lineas lata. Cymæ apice tricephalæ vel sæpius subumbellatim polycephalæ. Capsula 4 lineas diam. lata. Inter affines inflorescentiâ et capsulâ tuberculatâ notabilis. (v. s.)

β, *Canariensis*, foliis abbreviatis paniculâ subbrevioribus nec eam superantibus, capsulâ vix piso majore, semine obsolete carinato carunculâ patellari ovatâ. — In sylvaticis Teneriffæ ad Taganana (Bourgeau n. 1515!), prope Anaga (de la Perraudière!), in insulâ Palmâ (Bourgeau n. 976!). Arbor sæpe 30-pedalis. (v. s.)

420. *E. STYGIANA* (Watson in Hook. Lond. Journ. III, p. 605), fruticosa, ramis erectis superne dense foliosis, cymis pedunculatis apice 3-4-fidis bifidis paniculam terminalem corymbosam formantibus junioribus hirtis, foliis subsessilibus e basi vix attenuatâ lineari-oblongis subtus glaucis nervo medio manifesto percursis vix venulosis sparsim pilosulis, floralibus oblongis concaviusculis villosis, involucri extus pilosuli intus dense hirsuti lobis ovatis, glandulis semiorbiculatis, capsulâ verrucosâ ... ♂ In montibus insul. Azoricarum Fayal et Flores alt. 1500'-2500' (Watson!). *E. mellifera* Seub. fl. Azor. n. 172, non Ait. Valde affinis præcedenti et forsan ejus varietas, differre videtur foliis brevioribus non venulosis nervo latiori percursis, basi vix attenuatis, apice non longe mucronatis, partium floralium indumento. (v. s. spec. jun. in h. Kew.)

421. *E. ATROPURPUREA* (Brouss. el. hort. bot. Monsp. 1805, p. 24), fruticosa, glabra, dichotome ramosa, ramis lucidis denudatis apice dense et rosulatum foliosis, umbellæ 5-10 radiis 3-5-fidis, foliis e basi sessili attenuatâ lanceolatis vel obovato-lanceolatis obtusis, umbellaribus conformibus, floralibus sub involucri atropurpureis rarius luteis e basi breviter connatâ transverse ovatis obtusis vel retusis, involucri breviter campanulati extus et intus sub glandulis breviter hirtuli lobis ovatis ciliato-fimbriatis, glandulis 4 transverse ovatis, stylis brevibus ad medium connatis apice bifidis incrassatis, capsulæ ovatæ trilocæ coccis dorso rotundatis, semine oblongo-tetragono rugoso-scribiculo, carunculâ conicâ facie interiori sulcatâ. ♂ In rupestribus meridionalibus Teneriffæ (Bourgeau n. 471! et 1516!). Vulgo *Tabayba Majorera*. — Bot. mag. t. 3321, Webb et Berth. phyt. Canar. III, t. 207 et Tab. phyt. t. 2. Folia 2-3-pollicaria, 6-8 lineas lata, umbellâ sublongiora. Capsula magn. pisi. (v. s.)

422. *E. BOURGÆANA* (J. Gay in Bourg. pl. Canar. exs. 1855), fruticosa, glabra, ramis denudatis apice dense foliosis, umbellæ 4-6-fidæ radiis 2-3-fidis, foliis e basi attenuatâ late lineari-lanceolatis apice retusis, floralibus magnis flavidis vel purpurascensibus e basi breviter connatâ transverse ovatis retusis mucronulatis, involucri cylindrico-campanulati lobis oblongis truncato-retusis hirtis, glandulis 4 vel 5 minutis transverse oblongis truncatis breviter bicornutis, stylis alte connatis incrassato-bilobis, capsulæ profunde sulcatæ coccis dorso rotundatis, semine ovato profunde scribiculo-rugoso, carunculâ ovatâ cristæformi. ♂ In præruptis Teneriffæ ad Guimar (Bourgeau n. 1511!, de la Perraudière!). Foliis et inflorescentiâ affinis *E. atropurpureæ*, sed folia retusa, floralia multo ampliora, glandulæ bicornutæ, capsula duplo major 4 lin. lata profunde sulcata, etc. (v. s.)

423. *E. PISCATORIA* (Ait. Hort. Kew. ed. I, vol. II, p. 137), fruticosa, glabra, ramis parte superiori densiuscule foliosis, umbellæ radiis 8-12 brevibus breviter bifidis,

[Begin Page: Page 108]

108 DICOTYL. szv , EXOGENJE£. EUPHORBIA.

et 1510 !). Vulgo Tabayba dulce. — Webb phyt. Can. t. 200 et tab. phytost. t. 6. In Senegambi:e regno Cayor ad sepes formandas cultà sub nomine Selane (Heu-"delot!), an ibi vere spontanea ? Frutex 4-5-pedalis. Folia 9-12 lineas longa, 2-2 lata. Involucrã. quzedam hermaphrodita, siepius mascula. Capsula piso major. Suceus mitis et jucundus. Species a sequentibus inflorescentiã 1nonocephalã et defectu foliorum floralium distinctissima. (v. s.)

119. E. MELLIFERA (Ait. hort. Kew. ed. I, vol. 3, p. 403), glabra, arborescens, ramis superne foliosis, cymis pedunculatis paniculam terminalem simplicem vel compositam basi tantum foliosam formantibus, foliis basi' attenuatis elongato-lanceolatis acutis mucronatis subsessilibus nervis secundariis manifestis, foliis floralibus ad basin involucrorum biris oblongis concavis mucronatis ciliatis, involucri campanulati lobis late et breviter ovatis ciliatis, glandulis quinis semiorbiculatis subtus hirtulis, stylis longis ad medium coalitis bifidis apice capitatis, capsule 1nagze spherieze subtrisule:e sparsim tuberculatis coccis dorso rotundatis, semine ovato levi dorso carinato, carunculã patellari semilunatã. 5 In sylvaticis Madere (Webb! Mason!!). — Vent. Malm. t. 30,-Bot. mag. tab. 1305. E.longifolia Lam. dict. II, p. 417. Arbuscula 6-15-pedalis. Folia 6-12 poll. longa, 10-15 lineas lata. Cymse apice iricephale vel sepius subumbellatim polycephalee. Capsula 4 lineas diam. lata. Inter affines inflorescentiã et capsulã tuberculatã-notabilis. (v. s.)

DB. Canãriensis, foliis abbreviatis paniculã subbrevioribus nec eam superantibus, capsulã vix piso majore, semine obsoletius carinato carunculã patellari ovatã.— In sylvaticis TeneriITa: ad Taganana (Bourgeau n.1515 I), prope Anaga (de là Perraudière !), in insulã Palmã (Bourgeau n. 976 !). Arbor siepe 30-pedalis. (v. s.)

420. E. SrYerANA (Watson in Hook. Lond. Journ. III, p. 605), fruticosa, ramis erectis superne dense foliosis, cymis pedunculatis apice 3-4-fidis bifidis paniculam terminalem corymbosam formantibus. junioribus hirtis, foliis subsessilibus e hasi vix attenuatã lineari-oblongis subtus glaucis nervo medio manifesto percursis vix venulosis sparsim pilosulis, floralibus oblongis concaviusculis villosis, involucri extus pilosuli intus dense hirsuti lobis ovatis, glandulis semiorbiculatis, eapsulã verrucosã ... In montibus insul. Azoricarum Fayal et Flores alt. 1500/-2500' (Watson!). E. mellifera Seub. fl. Azor. n. 172, non Ait. Valde affinis precedenti et forsan ejus varietas, differre videtur foliis brevioribus non venulosis nervo latiori percursis, basi vix attenuatis, apice non longe mucronatis, partium floralium indumento. (v. s. spec. jun. in h. Kew.)

421. E. ATROPURPUREA (Brouss. el. hort. bot. Monsp. 1805, p. 24J, (ruticosa, glabra, dichotome ramosa, ramis lucidis denudatis apice dense et rosulatim foliosis, umbelli? 5-10 radiis 3-5-fidis, foliis e basi sessili attenuatã lanceolatis vel obovato-Janceolatis obtusis, umbellaribus conformibus , floralibus sub involucro atropurpureis rarius luteis e basi breviter connatã transverse ovatis obtusis vel retusis, involucri breviter campanulati extus et intus sub glandulis breviter hirtuli lobis ovatis ciliato-fimbriatis, glandulis 4 transverse ovatis, stylis brevibus ad medium eonnalis apice bifidis incrassatis, capsulie ovatze trisuleze coccis dorso rotundatis, semine oblongo-tetragono rugoso-scribiculato, carunculã conicã facie interiori sulcatã.. 5 In rupestribus meridionalibus Teneriffe (Bourgeau n. 471! et 1516 !). Vulgo Tabayba Majorera. — Bot. mag. t. 3321, Webb et Berth. phyt. Canar. III, t. 207 et Tab. phyt. t. 2. Folia 2-3-pollicaria, 6-8 lineas lata, umbellã sublongiora. Capsula magn. pisi. (v. s.)

422. E. BouRncEANA (J. Gay in Bourg. pl. Canar. exs. 1855), fruticosa, glabra, ramis .denudatis apice dense foliosis, umbellie 4-6-fidze radiis 9-3-fidis, foliis e basi at-

tenuatá late lineari-lanceolatis apice retusis, floralibus magnis flavidis vel pur-
purascenscentibus e basi breviter connatá transverse ovatis retusis mucronulatis,
involucris cylindrico-campanulatis lobis oblongis truncato-retusis hirtis, glandulis
4-5 minutis transverse oblongis truncatis breviter bicornulis, stylis alte con-
natis incrassato-bilobis, capsula: profunde sulcatá: e coccis dorso rotundatis, semine
ovato profunde scrobiculato-rugoso, carunculá ovatá cristiformi. 5 In prae-
Teneriffae ad Guimar (Bourgeau n. 1511!, de la Perraudière!). Foliis et inflore-
scentiá affinis E. atropurpurea, sed folia retusa, floralia multo ampliora, glandulis
bicornutis, capsula duplo major 4 lin. lata profunde sulcata, etc. (v. s.)
198. E. PISCARIA (Ait. Hort. Kew. ed. I, vol. II, p. 137), fruticosa, glabra, ramis
parte superiori densius foliosis, umbellae radiis 8-12 brevibus breviter bifidis,