



<https://www.biodiversitylibrary.org/>

**Species plantarum : exhibentes plantas rite cognitatas ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas**

Berlin, Junk, 1908

<https://www.biodiversitylibrary.org/bibliography/37656>

**t.1 (1753):** <https://www.biodiversitylibrary.org/item/84235>

Page(s): Page 450

Holding Institution: NCSU Libraries

Sponsored by: NCSU Libraries

Generated 25 March 2024 12:36 PM

<https://www.biodiversitylibrary.org/pdf4/1684848i00084235.pdf>

This page intentionally left blank.

Reseda floribus trigynis, calycibus maximis hexaphyllis.  
*Hort. ups.* 150.

Reseda minor vulgaris. *Tournef. inst.* 423.

Resedæ affinis Phyteuma. *Baub. pin.* 100. *prodr.* 42.  
*t.* 42.

Erucago apula trifida & quinquefolia. *Col. ecphr.* 267.  
*t.* 269.

β. Reseda minor vulgaris, folio minus inciso. *Tournef.*  
*inst.* 423.

γ. Reseda minor vulgaris, foliis integris. *Tournef. inst.* 423.  
Phyteuma. *Baub. hist.* 3. *p.* 386. *fig.*

*Habitat in Gallia, Italia.* ☉

## EUPHORBIA.

### \* *Fruticosa aculeata.*

*antiquorum.* 1. EUPHORBIA aculeata subnuda triangularis articula-  
ta: ramis patentibus. *Hort. cliff.* 196. \* *Hort. ups.*  
138. *Fl. zeyl.* 199. *Mat. med.* 154. *Roy. lugdb.*  
194. *Diss. euph.* 1. \*

Euphorbium antiquorum verum. *Comm. hort.* 1 *p.* 23.  
*t.* 12.

Schadidacalli. *Rheed. mal.* 2. *p.* 81. *t.* 42.

β. Tithymalus aizoides triangularis & quadrangularis ar-  
ticulofus & spinosus, ramis compressis. *Comm. præl.*  
21. *t.* 55. *f.* 5.

*Habitat in India.* †

*canariensis* 2. EUPHORBIA aculeata nuda subquingularis, a-  
culeis geminatis. *Hort. cliff.* 196. \* *Hort. ups.* 138.  
*Roy. lugdb.* 194. *Diss. euph.* 2. \*

Tithymalus aizoides fruticosus canariensis aphyllus qua-  
drangularis & quingularis, spinis geminis adun-  
cis atronitentibus armatus. *Comm. hort.* 2. *p.* 207  
*t.* 104.

Tithymalus aizoides lactifluus s. Euphorbia canariensis  
quadrilatera & quingulatera, cerei effigie. *Pluk. alm.*  
370. *t.* 320. *f.* 2.

*Habitat in Canariis.* †

*heptagona.* 3. EUPHORBIA aculeata nuda septemangularis: spinis  
solitariis subulatis floriferis. *Hort. cliff.* 196. \* *Roy.*  
*lugdb.* 194. *Diss. euph.* 6. \*

Euphorbium heptagonum, spinis longissimis in apice fru-  
giferis. *Bærb. lugdb.* 1 *p.* 258. *t.* 258.

*Habitat - - -* †

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 450]

45-0 DGDECANDRIA TRIGYNJA,

Reieda floribus trigynis, calycibus maxiniishexaphyllis

Hort. tipf. 15-0.

Rcfeda minor vulgaris. 'Tournef. infl. 423.

Refedae affinis Phyteuma. Btmh. fin. 100. frodr. 4^.

t, 42.

Erucago apula trifida & quinquefoUa. Col. ecphr.z6^.

t. 269.

/3. Refeda minor vulgaris, folio ininus incifo. T^ournef.

infi. 423.

y.Refedaminorvulgaris,foliis integris. ToHrnef. infl.^lIs

Phyteuma. Bauh. hifi. 3. />. 386. fig,

Habitat in Gallia, Italia. ©

EUPHORBJA.

\* Fruticeja aculeattc»

mfi^mrnm.i^ EUPHOREIA aculeata fubnuda triangularis articula-

ta: ramis pateutibus. Hart. chff. 796. \* Hort. npf.

138. fl. zeyl. 199. Mat. med.. 15-4. Roy. lugdh-

194. Dift\ ev.ph. I. \*

Euphorbium antiquorum verum. Comm. hort. i p. 23.

t. 12.

Schadidacalli. Rheed. mal. 2. p. 81. t. 42.

[3. Tithymalus aizoides triangularis & quadrangularis ar-

ticulofus & fpinofus, ramis comprelfis. Comm. pral

11. t. 3-5-. / 5-.

Hahitat in India. "b

rmarijtyp 2. EUPHORBIA aculeata nuda frbquingularis, a-

culeis geminatis. Hort. cUff. 196. \* fiort. upf. 138.

/?oy. lugdh. 194. Dift', ettph. 2. \*

Tithymalus aioides fruticosus canariejifis aphyllusqua'

drangularis & quingularis , fpinis geminis adun-

cis atronitentibus armatus. Comm. hort. z. p. 207

t. V104.

Tiihymalus aizoides laftifluus f. Euphorbia canarienfis

quadrilatera & quinquelatera, cereieffigie. Phk. alm.

370. t. 320. / . 1.

Hahitat in Canariis. t>

htftngtna. 3. EUPHORBIA aculeata nuda feptemangularis: fphns

ibliLariis fubulatis rloiiferis. Hort. clijf. 196. \* Rny.

hgdh. 794. Diff. eitph. 6. \*

Eupnorbiunheptagonum,fpinislongiffimis in apice fni-

giferis. Bocrh. lit^db. i ' p. ^jS. t. i-^i.

Hahitat - - - - t> I