



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789
<https://www.biodiversitylibrary.org/bibliography/4504>

v.1: <https://www.biodiversitylibrary.org/item/23432>

Page(s): Page 130

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 26 March 2024 11:28 AM
<https://www.biodiversitylibrary.org/pdf4/1685176i00023432.pdf>

This page intentionally left blank.

birta.

22. P. foliis ovatis glabris, floribus lateralibus. *Thunb. Protea, n. 55. Syst. veget. 141.*

Hairy Protea.

*Nat. of the Cape of Good Hope.**Introd. 1774, by Mr. Francis Masson.**Fl.*

G. H. h.

pubera.

23. P. foliis ovatis capitulisque terminalibus tomentosis. *Thunb. Protea, n. 56. Syst. veget. 141.*

Downy-leav'd Protea.

*Nat. of the Cape of Good Hope.**Introd. 1774, by Mr. Francis Masson.**Fl.*

G. H. h.

***** Rotundatæ: foliis subrotundis.

*cyna-
roides.*

24. P. foliis subrotundis petiolatis glabris. *Thunb. Protea, n. 59. Syst. veget. 142.*

Round-leav'd Protea.

*Nat. of the Cape of Good Hope.**Introd. 1774, by Mr. Francis Masson.**Fl.*

G. H. h.

GLOBULARIA. *Gen. pl. 112.*

Cal. communis imbricatus; proprius infundibuliformis, inferus. Recept. paleaceum.

longifolia.

1. G. caule fruticoso, foliis omnibus lineari-lanceolatis integerrimis, capitulis axillaribus.

Alypum five Herba terribilis procerior, cortice cinereo scabro, folio acuminato longiore. *Sloan.*

jam. 124. hist. 1. p. 19. t. 5. f. 3.

Long-leav'd Globularia.

*Nat. of Madeira.**Introd. 1775, by Sir Joseph Banks, Bart.**Fl. July and August.*

G. H. h.

2. G.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 130]

150

birta.

- pubera,

çyna-

roides.

ee tifa.

=s G. caule fruticofo, foliis omnibus Losicns a

@y

TETRANDRIA. re eer ~ Protea,

Sa°P; foliis ovatis glabris, floribus lateralibus,, Thin

Protea, nm. 55... Syft. veget. stein

Hairy Protea,

_ Nat. of the Cape of Good Hope. Lite

Introd. 1774 the Mr. Francis Mafion, | its

23. P, foliis ovatis capitulifque terminalibus. se

Thunb. Protea, n. 56. Syft. veget. is rae

Downy-leav'd Protea,

Nat. of the Cape of Good Hope. . iE ne

Introd: 1774 by Mr. Francis Maffon, _ A

\ G Hh

eR eRS Bideciies: foliis febrethlle n

24. P. foliis fubrotundis petiolatis glabris. Thank Prt,

n. 59. Syft. veget. 142.

Round-leav'd Protea.

Nat. of the Cape of Good Hope.

Introd. 1774, ar Mr. Francis Maffon.

FI 4

«~ GLOBULARIA. Gem pm om

_ Gal. commanis imbricatus ; proprius infun

inferus., Recept. paleaceum. 4

integerrimis, capitulis axillaribus.

Alypum five Herba terribilis procerior, ç

hereo ftabro, folio acuminato longiore.

jam. 124. bift. 1. p.19. t. 5. fe % :

Long-leav'd Globularia. |

Nat. of Madeira,

pte 1775, by Sir Jofeph Ranks, Bart,

Fi. July and ADgUR,