



<https://www.biodiversitylibrary.org/>

The genera and species of orchidaceous plants

London, Ridgways, 1830-1840

<https://www.biodiversitylibrary.org/bibliography/499>

Item: <https://www.biodiversitylibrary.org/item/9889>

Page(s): Page 318

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 27 March 2024 11:44 AM

<https://www.biodiversitylibrary.org/pdf4/1685625i00009889.pdf>

This page intentionally left blank.

longioribus, labelli tripartiti laciniis filiformibus lateralibus recurvis brevioribus, calcare pendulo filiformi ovario longiore.

Orchid. arenaria. Burch. Hb. 5654.

Hab. ad Caput B. Spei, Burchell. (hab. s. sp. comm. cel. Burchell.)

Caulis 1-1½-pedalis. Ovaria abruptè pedicellata. Sepala lateralia pendula.

53. HABENARIA filicornis.

H. caule gracillimo bifolio, foliis strictissimis vaginantibus oblongo-linearibus obtusis, racemo laxo multifloro, bracteis subulatis pedicellorum longitudine, labelli tripartiti laciniis filiformibus obtusis intermediâ ascendentे lateralibus deflexis duplò longiore, calcare filiformi compresso ovarii longitudine.

Orchis filicornis. Thonning! in Schumach. Beskriv. af Guineisk. Plant. 2, 171.

Hab. in Guinea, Thonning. (exam. s. sp. in hb. Hornem.)

Caulis 1-1½-pedalis. Foliorum lamina caule parallela obtusa. Flores minuti, ichneumonis facie. Petala falcata cum sepalo supr. refracto in galea agglutinata. Sepala lateralia majora retrorsum falcata. Anthera acuminata; basibus solutis brevibus erectis. Processus carnosus acutus decurvi.

54. HABENARIA procera.

H. "labello tripartito : laciniis ascendentibus lateralibus linearibus divaricatis mediâ subulatâ porrectâ canaliculatâ, calcare linearis obtuso germine longiore, foliolis cal. lateralibus exterioribus deltoideis."

Orchis procera. Swartz in Pers. Syn. 1, 506.

Hab. in Sierra Leone, Afzelius.

55. HABENARIA tridactylites.

H. foliis 2 radicalibus oblongo-lanceolatis petiolatis, racemo laxo paucifloro, bracteis membranaceis acuminatis ovario duplò brevioribus, labelli tripartiti laciniis filiformibus intermediâ paulò longioribus, calcare filiformi clavato pendulo ovarii longitudine.

Orchis tridactylites. Webb et Berth. Fl. Can. ined.

Hab. ad rupes, et in cæspitosis ins. Canariensis frequens, Webb et Berthellot. (hab. s. sp. comm. cell. W. et B.)

Caulis spithameus v. minor, foliis radicalibus paulò altior. Sepalum supremum lato-ovatum acutum cum petalis angustis falcatis galeam planiusculum efformans. Sepala lateralia linearia obtusiuscula. Anthera lata, lobis parùm solutis divaricatis; processus carnosus oblongi distantes.

56. HABENARIA graminea.

H. foliis radicalibus linearibus acuminatis scapo filiformi stricto multò brevioribus, racemo laxo paucifloro, bracteis squamiformibus acutis vix pedicello ovarii longioribus, labelli tripartiti sepalorum longitudine laciniis filiformibus subæqualibus, calcare compresso clavato ovarii longitudine.

Platanthera linifolia. Lindl. in Wall. Cat. no. 7041.

Hab. in montibus Silhet, Wallich. (hab. s. sp. e Mus. Angl. Ind.)

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 318]

318 OPHRYDE.;

longioribus, labelli tripartiti laciniis filiformibus lateralibus recurvis

brevioribus, calcare pendulo filiformi ovario longiore.

Orchid. arenaria. Burch. Hb. 5654.

Hab. ad Caput B. Spei, Burchell. (hab. s. sp. comm. cel. Burchell.)

Caulis 1-14-pedalis. Ovaria abrupte pedicellata, Sepala lateralia pendula.

53. Hapenaria filicornis.

H. caule gracillimo bifolio, foliis strictissimis vaginantibus oblongo-linearibus obtusis, racemo laxo multifloro, bracteis subulatis pedicellorum longitudine, labelli tripartiti laciniis filiformibus obtusis intermedia ascendentे lateralibus deflexis duplo longiore, calcare filiformi compresso ovarii longitudine.

Orchis filicornis. Zhonning ! in Schumach. Beskriv. af Guineisk. Plant. 2, 171.

Hab. in Guinea, Thonning. (exam. s. sp. in hb. Hornem.)

Caulis 1-li-pedalis. Foliorum lamina caule parallela obtusa. Flores minuti, ichneumonis facie. Petala falcata cum sepalo supr. refracto in galea agglutinata. Sepala lateralia majora retrorsum falcata. Anthera acuminata; basibus solutis brevibus erectis. Processus carnosus acutus decurvi.

'54, HaBENARIA procera.

H. " labello tripartito : laciniis ascendentibus lateralibus linearibus divaricatis mediа subulata porrectа canaliculatа, calcare linearis obtuso germine longiore, foliolis cal. lateralibus exterioribus deltoideis. '

*Orchis procera. Swartz in Pers. Syn. 1, 506.

'Hab. in Sierra Leone, Afzelius.

55. Hapenaria tridactylites.

H. foliis 2 radicalibus oblongo-lanceolatis petiolatis, racemo laxo lwh floro, bracteis membranaceis acuminatis ovario duplo brevioribus, labe tripartiti laciniis filiformibus intermediа paulo longioribus, calcare filiformi clavato pendulo ovarii longitudine.

"Orchis tridactylites. Webb et Berth. Fl. Can. ined.

'Hab. ad rupes, et in caspitosis ins. Canariensium frequens, Webb et -Berthellot. (hab. s. sp. comm. cell. W. et B.)

'Caulis spithameus v. minor, foliis radicalibus paulo altior. | gione supremum lato-ovatum acutum cum petalis angustis falcatis usque planiusculum eiformans. Sepala lateralia linearia obtusiuscula. i thera lata, lobis parum solutis divaricatis; processus carnosus oblong distantes.

56. HABENARIA graminea.

H. foliis radicalibus linearibus acuminatis scapo filiformi stricto pup brevioribus, racemo laxo paucifloro, bracteis squamiformibus usque pedicello ovarii longioribus, labelli tripartiti sepalorum longitudo! ii laciniis filiformibus subaequalibus, calcare compresso clavato ovarii longitudine.

Platanthera linifolia. Lindl. in Wall. Cat. no. 7041.

Hab. in montibus Silhet, Wallich. (hab. s. sp. e Mus. Angl. Ind.)