



<https://www.biodiversitylibrary.org/>

**Synopsis plantarum succulentarum :cum descriptionibus,
synonymis, locis, observationibus anglicanis, culturaque**

Londini, Typis R. Taylor et Socii, 1812

<https://www.biodiversitylibrary.org/bibliography/9462>

Item: <https://www.biodiversitylibrary.org/item/38208>

Page(s): Page 312, Page 313

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 31 March 2024 11:36 AM

<https://www.biodiversitylibrary.org/pdf4/1686535i00038208.pdf>

This page intentionally left blank.

HABITAT C. B. S.

S. h.

FLORET

INTR. D. 1790.

OBS. I have seen this remarkable plant alive in the matchless gardens of Kew; but it is not at present to be found there, or elsewhere in Britain.

Classis XIX.

SYNGENESIA

ÆQUALIS.

KLEINIA. (*Linn. in hort. cliff.*)—*Cacalia*
Linn. &c.

Flores discoidei. Calyx cylindricus, basi tan-
tum subbracteolatus. Receptaculum nudum.
Pappus pilosus.

OBS. *Caulis fruticosus, foliaque succulenta*
glauca, intus resinosa.

Nereifo- K. (Oleander-leav'd) caule fruticoso, foliis lan-

ceolatis planis, floribus corymbosis. Willd.

1. sp. pl. 3. 726. (*Cacalia Kleinia.*)—*Kleinia,*
foliis lanceolatis planis, caule lœvi ventricoso.

Linn. hort. cliff. 395.—Plant. grass. cum
icone.—Cacalianthemum folio nerii glauco.

Dill. elth. 61. t. 54. f. 2.

HABITAT in Canariis. G. H. h.

FLORET Sept. Oct.

CULT. 1732.

OBS. The original Linnaean name of Kleinia ought to be restored to this plant and its allies. But I am not certain the following ought to associate with it, as some of them have calyxes much more numerously divided.—Cacalia, as it at present stands, is an heterogeneous mass.

K. (Rugged) caule fruticoso papillis cylindra- *papilla-*
ceis truncatis petiolaribus obsito, foliis lance- *ris.*
olatis planis. Willd. sp. pl. 3. p. 1725.— 2.
(*Cacalia papillaris*) *Kleinia caule carnosum,*
petiolis truncatis obvallato. Linn. hort. cliff.
395.—*Cacalianthemum caudice papillari.* Dill.
elth. 63. t. 55. f. 63.

HABITAT C. B. S. G. H. h.

FLORET

CULT. 1727.

K. (Fig-marygold) caule fruticoso, foliis com- *Ficoides.*
pressis, carnosis. Willd. sp. pl. 3. p. 1727.— 3.
(*Cacalia Ficoides*).—*Kleinia foliis carnosis*
lanceolatis compressis, caule tereti. Linn. hort.
cliff. 395.—*Senecio africana arborescens: fi-*
coidis folio et facie. Comm. rar. 40. t. 40.—
Bradl. succ. 5. p. 11. t. 49.

HABITAT C. B. S. G. H. h.

FLORET Jun.—Nov.

CULT. 1727.

K. (Creeping-rooted) caule fruticoso, foliis de- *repens.*
pressis carnosis. Willd. sp. pl. 3. 1727.— 4.
(*Cacalia repens*) Thunb. prod. 141.—*Plant.*
grass. cum icono.

HABITAT C. B. S. G. H. h.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 312]

31g Monadelphia Heptand. | Monsonia.

HasnrraT C. B. S. Sh.

LORNFTI.:

I5TrR^D. 1790.

Ozs. I bave seen this remarkable plant alive in
the matchless gardens of Kew ; but it is notâ
present to befound tbere, or elsewhere i
Britain.

Classis XIX.

SYNGENESIA

ZQUALIS.

KLEINIA. (Linn. in hort. cliff.) —Cacalia

Linn. (3c,

Flores discoidei. Calyz cylindricus, basi tan-
tum NOROPIIMUR, Receptaculum. nudum.

Pappus pilosus.

Ons. Caulis wies foliaque | succulenta
auca, intus resinosa,

Me on K. (Oleander-leav'd) caule fruticoso, foliis lan-
lia, ccolatis planis, floribus corymbosis,
m pL. ST ae Tides Kleinia.) — Kieinia

Ju Jiémd nis, caule leevi ventricoso.

ipo ent p" ; z lgbcoilent, grass. cum
icone.— "i folio meri gie

Dill. elih. 61. 4. 54. rq 2. |

i

[Begin Page: Page 313]

Syngenesia /Equalis. Kleinia. —— 313

HanBrTaT in Canariis. : ' G.H. b.

FronzT Sept. Oct.

Curr. 1732.

Oss. The original Licitfftem name of Kleintia
ought to be restored to this plant and its allies.
But I am not certain the followifig ought to
associate with it, as some of them have ca-
lyxes much more numerously. divided.—Ca-
calia, as it at present stands, is an heterogene-
ous mass.

K. (Rugged) caule fruticoso papillis cylindra- papilla-
ceis truncatis petiolaribus obsito, foliis lance- ris.

olatis planis. JVlld. sp. pl. 3. p. 1725.— .— 2.

(Cacalia papillaris) Kleinia caule carnosso,
need tr uncatis. obvallato.. Linn. hort. cli ci
elil. 03. 655 J. 03.

HasrraT C. B. S. G. H. B.

FroskT.... 2

Curr. 1727.

K. (Fig-marygold) caule fruticoso, foliis com- Ficoides.
pressis, carnosis. Willd.sp. pl. 3. p. 1727.— . 3.

lia Ficoides). — Kleinia foliis carnosis
lanceolatis compressis, caule tereti. Linn. hort.
cliff. 395.— Senecio fe arborescens: fi-
coidis folio et facie. Comm. rar. 40. 1. 40.—

Bradl. succ. 5. J^ 11. £. 49.

HanrrAT C. B. S G.H. 5.

FrongT DdirsedoK:

Curr. 1727.

K. (Creeping-rooted) caule driicolo, foliis de- repens.
pressis carnosis. JVilld. sp. pl. 3. 1727.— . 4.

penes rd ina Thunb. prod. 141.— Plant.

Le

eee BS. JA de z G.H. 5.