



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789
<https://www.biodiversitylibrary.org/bibliography/4504>

v.3: <https://www.biodiversitylibrary.org/item/23434>

Page(s): Text, Title Page, Page 396, Page 397

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 27 April 2024 12:08 PM
<https://www.biodiversitylibrary.org/pdf4/1695203i00023434.pdf>

This page intentionally left blank.

G-73
K48
178
V.
HORTUS KEWENSIS;

O R, A
C A T A L O G U E
O F T H E
P L A N T S
CULTIVATED IN THE
ROYAL BOTANIC GARDEN AT KEW.

B Y W I L L I A M A I T O N,
GARDENER TO
H I S M A J E S T Y.

I N T H R E E V O L U M E S.

V O L. III.

DIADELPHIA — CRYPTOGAMIA.

L O N D O N:

PRINTED FOR GEORGE NICOL, BOOKSELLER
TO HIS MAJESTY, PALL MALL.

M.DCC.LXXXIX.

Introd. 1774, by Mr. Francis Masson.

Fl. July.

G. H. h.

HIPPOPHAE. *Gen. pl.* 1106.

MASC. *Cal.* 2-partitus. *Cor.* o.

FEM. *Cal.* 2-fidus. *Cor.* o. *Styl.* 1. *Bacca* mono-sperma.

Rham-noides. 1. *H.* foliis lanceolatis. *Sp. pl.* 1452.

Common Sea Buck-thorn.

Nat. of England,

Fl. April.

H. h.

MYRICA. *Gen. pl.* 1107.

MASC. *Amenti* squama lunata. *Cor.* o.

FEM. *Amenti* squama lunata. *Cor.* o. *Styli* 2. *Bacca* 1-sperma.

Gale. 1. *M.* foliis lanceolatis subserratis, caule suffruticoso. *Sp. pl.* 1453.

Sweet Gale, Willow, or Candleberry Myrtle.

Nat. of Britain.

Fl. May.

H. h.

cerifera. 2. *M.* foliis lanceolatis subserratis, caule arborescente. *Sp. pl.* 1453.

angusti-folia. α *Myrtus* brabanticae similis carolinensis baccata, fructu racemoso sessili monopyreno. *Catesb. car.* I. p. 69. t. 69.

Common American Candleberry Myrtle.

latifolia. β *Myrtus* brabanticae similis carolinensis humilior, foliis latioribus et magis ferratis. *Catesb. car.* I. p. 13. t. 13.

Broad-leav'd American Candleberry Myrtle.

Nat.

Nat. of North America.

Cult. 1699, by the Dutches of Beaufort. *Br. Mus.*

Sloan. mss. 525 & 3349.

Fl. May and June.

H. b.

3. *M. foliis elliptico-lanceolatis subserratis, amentis mas-* *Faya.*
culis compositis, drupa nucleo quadriloculari.

Azorian Candleberry Myrtle.

Nat. of Madeira and the Azores. *Mr. Fr. Masson.*

Introd. 1777.

Fl. June and July.

H. b.

4. *M. foliis oblongis opposite sinuatis. Sp. pl. 1453.* *quercifo-*
lia.
α Myrica quercifolia, foliis oblongis opposite sinuatis gla-
bris. Mill. dict.

Smooth Oak-leav'd Candleberry Myrtle.

β Myrica birsuta, foliis oblongis opposite sinuatis hirfu-
tis. Mill. dict.

Hairy Oak-leav'd Candleberry Myrtle.

Nat. of the Cape of Good Hope.

Cult. 1752, by Mr. Ph. Miller. *Mill. dict. edit. 6. n. 5.*

Fl. June and July.

G. H. b.

5. *M. foliis subcordatis serratis sessilibus. Sp. pl. 1454.* *cordifolia.*

Heart-leav'd Candleberry Myrtle.

Nat. of the Cape of Good Hope.

Cult. 1759, by Mr. Ph. Miller. *Mill. dict. edit. 7. n. 7.*

Fl. May and June.

G. H. b.

B R U C E A. L'Herit. stirp. nov.

MASC. *Cal. 4-phyllo. Petala 4.*

FEM. *Cal. et Cor. ut in mare. Peric. 4, monosperma.*

1. BRUCEA. *L'Herit. stirp. nov. p. 19. tab. 10.*

Brucea antidyserterica. J. F. Miller. ic. tab. 25.

ferrugi-
nea.

African

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Text, Title Page]

Pk fe
HORTUS KEWENSIS;
ie eer
ora t A Ae Ot OU
OF ok ae
rE L A N ee S
CULTIVATED IN THE
ROYAL BOTANIC GARDEN AT KEW,
S2-PMTLLIAM AITON,
GARDENER TO
ELIS MAT BST Y.
IN THREE VOLUMES,
Ve dae
DIADELPHIA —CRYPTOGAMIA,
L O.N.D.O WN;
FRINTED FOR GEORGE NICOL, BOOKSELLER »
JO HIS MAJESTY, PALL MALL.
M.DCC,LXXXIX,
ee, ,

[Begin Page: Page 396]

cana ee, mere mecacnm can ieee
Jatifolia, :
ear
396
Rham-
noides. —
Gale.
cerifera,
_angufti-
folia,
DIOECIA TETRANDRIA,
Introd. 1774, by Mr. Francis Maffon.
Fi, July. G. Hb.
HIPPOPHAE. Gen. ph 1106,
Masc. Cal. 2-partitus. Cor. o.
Fem. Cal, 2-fidus, Cor. o. Styl 3, Bacca mono-
{perma. e.
1. H. foliis lanceolatis. Sp. pl. 1452.
Common Sea Buck-thorn.
Nat. of England,
Fi. April. H. hs
MY RICA. Gen. pl. 1107,
Masc. Amenti fquama lunata. - Gor. o.
Fem. Amenti fquama lunata. Cor. o. Styli 2, Bacca
1-{perma.
1. M. foliis lanceolatis fubferratis, caule fuffruticofo. Sp.
pl. 1453.

Sweet Gale, Willow, or Candleberry Myrtle.
Nat. of Britain,
Fi. May. H. k-
2. M. foliis lanceolatis fubferratis, caule arborefcente.
Sp. pl. 1453.
a Myrtus brabantice fimilis carolinensis baccata,
fructu racemofo feffili monopyreno, Catefh. cars 1s
p. 69. t. 69. :
Common American Candleberry Myrtle.
& Myrtus. brabantice fimilis carolinensis humilior,
foliis latioribus et magis ferratis. Catefb. car. Ts
p-13-% 13.
Broad-leav'd American Gindlebersy Myrtle. "
at,

[Begin Page: Page 397]

DIOECIA TETRANDRIA. Myrica. 397
Nat. of North America. : sed
Cult. 1699, by the Dutches of Beaufort. Br. Muf.
Shan. mff. 525 & 3349. |
F}, May and June, ae oars 1s
4. M. foliis elliptico-lanceolatis fubferratis, amentis maf- Faya,
4.
a
ns.
I,
culis compofitis, drupa nucleo quadriloculari.
Azorian Candleberry Myrtle. ee
Nat. of Madeira and the Azores. Mr. Fr. Maffon.
Introgeiqzy. 2 aoe Se
Fi], June and July. H. h.
M. foliis oblongis oppofite finuatis. Sp. pl 1453. quercifor
Myrica guercifolia, foliis oblongis Snponte finuatis gla- "4.
bris. Gil. dit... ~
Smooth Oak-leav'd Candleberry Myrtle.
Myrica bifluta, foliis oblongis. pppofite finuatis hirfu-
tis. AGI). did?. =errelit
Hairy Oak-leav'd Candleberry Myrtle.
Nat. of the Cape of Good Hope.
Cult. 1752, by Mr. Ph. Miller. AGI, idk. edit. 6. ms 5.
Fi, June and July. e 'G. Hi. h..
M. foliis subcordatis ferratis feffilibus. Sp. p.1454. cordifolia.
Heart-leav'd Candleberry Myrtle.
Nat. of the Cape of Good Hope.
Cult. 1759, by Mr. Ph. Miller. AGU, did. edit. 7. nN. 76
Fi, May and June, G.. He %..
BRUCEA. L'Herit. ftirg: any'
Masc. Cal. 4-phylus. Petala 4. ,
Fem. Cal. et Gor. ut in mare. Peric. 4, monofperma.
Brucea. L'Herit. flirp. nov. p. 19. tab. 10. _ ferrugia
Brucea antidyfenterica. F. F. Miller. ic. tab.25. nea.
African "