



<https://www.biodiversitylibrary.org/>

**Flora atlantica :sive historia plantarum quae in Atlante,
agro tunetano et algeriensi crescunt**

Parisiis, L.G. Desgranges, [1798-1799?]

<https://www.biodiversitylibrary.org/bibliography/323>

v. 2: <https://www.biodiversitylibrary.org/item/7542>

Page(s): Text, Title Page, Page 147

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 29 September 2024 6:21 AM

<https://www.biodiversitylibrary.org/pdf4/1738952i00007542.pdf>

This page intentionally left blank.

FLORA ATLANTICA,

S I V E

HISTORIA PLANTARUM,

QUÆ IN ATLANTE, AGRO TUNETANO

ET ALGERIENSI CRESCUNT.

AUCTORE RENATO DESFONTAINES, Instituti
nationalis Scientiarum Galliae socio, necnon in Museo
Historiae naturalis Parisiensi Botanices professore.

TOMUS SECUNDUS.

acc. to Soc Biol
Nat. Hist. Soc. 1: 148,
dates of fascicles of Vol
2 as follows:

-
- Livr. 5 - pp. 1-80, pls. 121-150, Oct. 1798.
6 - 81 - "Diadelphica Decandra" pls. 151-180, Nov. 1798.
7 - "Diadelphica Decandra" - "Polyadelphica Decandra", pls. 181-210, Feb. 1799.
8 - "Polyadelphica Decandra" - p. ? pls. 211-240, June 1799.
9 - pages ? - 458, pls. 241-261, July 1799.

PARISIIS,

Apud editorem L. G. DESGRANGES, in vico Haute-Feuille, N^o 14.

Typis C. PANCKOUCKE.

ANNO SEXTO REIPUBLICÆ GALLICÆ.

CAULIS erectus, ramosus, villosus, 2 — 3 decimetr. Folia petiolata, ternata, vix pubescentia; foliolis obovatis, denticulatis, 7—9 millimetr. latis, 9—11 longis. Stipulae suborbiculatae, dentatae, parvae, petiolo breviores. Flores laxi, axillares, solitarii, pedicellati; pedicellis filiformibus, folio longioribus, pubescentibus, 2 centimetr. longis. Calyx profunde quinquepartitus; laciniis lanceolato-subulatis, ciliatis. Corolla caerulea, longitudine calycis. Legumen nutans, exsertum, lineare, inflatum, villosum, 9—11 millimetr. longum, polyspermum.

HABITAT in collibus incultis Algeriae. ☉

ONONIS PENDULA. Tab. 191.

ONONIS foliolis ternis, obovatis, apice rotundatis; floribus racemosis, cernuis; pedicellis strictis; leguminibus villosis, calyce longioribus.

CAULIS villosus, erectus, 3 decimetr., ramosus. Folia petiolata, ternata; foliolis obovatis, glabris, superne dentatis, apice rotundatis, 9 millimetr. latis, 11 — 13 longis. Stipulae parvae, adpressae, ovatae, obtusae, denticulatae. Flores racemosi, pedicellati. Racemus 5 — 7 centimetr. Bractea foliacea, oblonga, simplices, apice dentatae. Pedicelli mutici, stricti, apice deflexi, bractea triplo breviores. Calyx quinquepartitus, villosus; laciniis lanceolato-subulatis, corolla paululum brevioribus. Corolla pallide caerulea. Flores absolutam fructificationem penduli. Legumen oblongum, lineare, acuminatum, inflatum, villosum, calyce paulo longius, polyspermum.

HABITAT prope Mascar in arvis. ☉

ONONIS VILLOSISSIMA. Tab. 192.

ONONIS villosissima, viscosa; foliis ternatis; floribus racemosis, confertis, sessilibus; corollis leguminibusque calyce brevioribus.

PLANTA villosissima et viscosa. Caulis herbaceus, ramosus, diffusus, 2—3 decimetr. Folia ternata, petiolata; foliolis obovatis aut ovato-rotundatis, denticulatis. Stipulae parvae, rotundatae, dentatae. Flores axillares, racemosi, conferti, solitarii, sessiles aut subsessiles. Calyx quinquepartitus; laciniis lanceolatis. Corolla parva, rosea, calyce brevior. Legumen ovatum, obtusum, compressum, villosum, calyce tectum, subdispermum.

HABITAT in arvis Algeriae.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Text, Title Page]

|
|
|

FLORA ATLANTICA;
Sirvr

HISTORIA. PLANTARUM,

QUA IN ATLA

TE, AGRO TUNETANO

ET ALGERIENSI :CRESCUNT

AucTORE RENATO DESFONTAINES , Instituti
nationalis Scientiarum Galliz: socio, necnon in Musco
Historic naturalis Parisiensi Botanices professore.

T.OMUS SECUNDUS, «^

Typs C. PANCKOUCK E.

ANNO SEXTO REIPUBLIGE GALLIQA.

[Begin Page: Page 147]

S.
imis 5g
dai, i.

,

pnis

g.

b

DIADELPHIA DECANDRIA. 147

CAULIS erectus, ramosus, villosus, 2 — 3 decimetr. Folia petiolata, ternata, vix pubescentia; foliolis obovatis, denticulatis, 7—9 millimetr. latis, 9—11 longis. Stipule: suborbiculate, dentate, parve, petiolo breviores. Flores laxi, axillares, solitarii, pedicellati; pedicellis filiformibus, folio longioribus, pubescentibus, 2 centimetr. longis. Calyx profunde quinquepartitus; laciniis lanceolato-subulatis, ciliatis. Corolla cocculescens, longitudine calycis. Legumen nutans, exsertum, lineare, inflatum, villosum, 9—11 millimetr. longum, polyspermum.

HABITAT in collibus incultis Algeriz. C)

ONONIS PENDULA. Tab. 1g.

floribus racemosis,
us villosis, calyce longioribus.

Oxoris foliolis teris, obovatis, ap-
cernuis; pedicellis stitucis; legu-

CAULIS villosus, erectus, 3 decimetr., ramosus. Folia petiolata, ternata foliolis obovatis, glabris, superne dentatis, apice rotundatis, 9 millimeu- latis, 11 — 13 longis. Stipule parva, adpresse, ova, obtuse, denti- culataz. Flores racemosi, pedicellati. Racemus 5— 7 centimetr. Bractez foliacez: oblonga, simplices, apice dentate. Pedicelli muici, stricti, apice deflexi, bractez wiplo breviores. Calyx quinquepartitus, villosus; laciniis lanceolato-subulatis, corollae paululum brevioribus. Corolla pallide cocculescens. Flores absolutae fructificatione penduli. Legumen oblongum, lineare, acuminatum, inflatum, villosum, . calyce paulo longius, polyspermum.

HABITAT prope Mascar in arvis.

ONONIS VILLOSISSIMA. Tab. 192.

Oxoxis villosissima, viscosa; foliis ternatis; floribus racemosis, confertis, sessilibus; corollis leguminibusque calyce brevioribus.

PLANTA villosissima et viscosa. Caulis herbaceus, ramosus, diffusus, 2— decimetr. Folia ternata, petiolata; foliolis obovatis aut ovato-rotun- datis, denticulatis. Stipula: parve, rotundataz, dentataz. Flores axillares, sacemosi, conferti, solitarii, sessiles aut subsessiles. Calyx quinqueparii- tus; laciniis lanceolatis: Corolla parva, rosea, calyce brevior. Legumen ovatum, obtusum, compressum, villosum, calyce tectum, subdispermum.

HABITAT in arvis Algeria.