



<https://www.biodiversitylibrary.org/>

**Caroli Linnaei ... Species plantarum :exhibentes plantas
rite cognitas, ad genera relatas, cum differentiis specificis,
nominibus trivialibus, synonymis selectis, locis natalibus,
secundum systema sexuale digestas**

Holmiae, Impensis Direct. Laurentii Salvii, 1762-63

<https://www.biodiversitylibrary.org/bibliography/11179>

v.2: <https://www.biodiversitylibrary.org/item/43462>

Page(s): Text, Title Page, Page 1011

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 29 September 2024 7:02 AM
<https://www.biodiversitylibrary.org/pdf4/1738963i00043462.pdf>

This page intentionally left blank.

Linnæan
QK91
C500
1762-63
v. 2

CAROLI LINNÆI

Equitis aur. de Stella polari,

ARCHIATRI REGII, MED. & BOTAN. PROFESS. UPSAL. ACAD.
PARIS. UPSAL. HOLMENS. PETROPOL. BEROL. IMPERIAL.
BERNENS. LONDIN. ANGLIC. EDENBURG. MONSPEL.
TOLOS. FLORENT. SOC.

SPECIES PLANTARUM,

EXHIBENTES
PLANTAS RITE COGNITAS,
AD

GENERA RELATAS,

CUM

DIFFERENTIIS SPECIFICIS,
NOMINIBUS TRIVIALIBUS,
SYNONYMIS SELECTIS,
LOCIS NATALIBUS,
SECUNDUM

SYSTEMA SEXUALE DIGESTAS.

MISSOURI
BOTANICAL
GARDEN.

TOMUS II.

Editio Secunda, aucta.

Cum Privilegio S. R M:tis Sveciae & S. R. M:tis Poloniae ac Electoris Saxon.

HOLMIAE
IMPENSIS DIRECT. LAURENTII SALVII
1763.

M.

Cicer sylvestre latifolium triphyllum. Baub. pin. 347.

Cicer sylvestre tertium. Dod. pempt. 525.

Cicer sylvestre verius. Lab. ic. 2. p. 73.

Habitat in Alpibus Helveticis, ḥ.

Vexillum pubescens. Ale carina breviores. Calyces cincti bracteis 3, cordatis, approximatis, brevioribus. Folia parva, integerrima, glabra.

* * * *

17. ONONIS spicis sessilibus, leguminibus nutantibus cernua, linearibus pilosis. Amoen. acad. 6. afr. 42. *.

Lotus africana annua hirsuta, floribus luteis. Comm. bort. 2. p. 163. t. 82.

Habitat ad Cap. b. spei. ○.

18. ONONIS racemis pedunculatis longis, foliis ternis capensis. tis suborbiculatis. Amoen. acad. 6. afr. 41. *.

Habitat ad Cap. b. spei. ○.

19. ONONIS pedunculis unifloris muticis, fructibus cernuis, foliis ternatis subrotundis crenatis.

Anonis pusilla hirsuta, flore ex albo & rubro colore mixto. Moris. hist. 2. p. 171.

Anonis purpurea non spinosa minor. Barr. ic. 354. mala.

Anonis annua pumila, flore purpurascente. Tournef. inst. 408.

Habitat in Delphinatu, Hispania, Italia. ○.

Planta viscosa-pubescentia, parva, diffusa, patula. Folia petiolata, ternata, orbiculata, crassifolia, plicata, crezata, viscosa-pubescentia. Stipulas ovales, patulae, obtuse, ferratae. Pedunculi axillares, uniflori, mutici, longitudine vix foliorum. Calyces longitudine pedunculi. Corollæ albidae vexillo subtus purpureo. Calyx defloratus reflectitur apice ad caulem. Legumen calyce sesquilongius, villosum.

ANTHYLLIS.

* Herbaceæ.

1. ANTHYLLIS herbacea, foliis quaterno-pinnatis, floribus lateralibus. Hort. upf. 221. * Saur. monsp. 237. tetraphylla.

Anthyllis foliis quinato-pinnatis inæqualibus, floribus lateralibus. Hort. cliff. 371. Roy. lugdb. 389.

Lotus pentaphyllum veticaria. Baub. pin. 332.

Trifolium halicacabum. Cam. bort. 171. t. 47.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Text, Title Page]

f

1

CAROLI LINKJBI

. iTti

Equitis aur. de Stella folari^ ^ 2L

^RCHIATRI ReGH, MeD & BoTAN. PROFESS. IJPSAL. AcAD.

Faris. Uj>sal. Holmens. Petropol Berol Imperial.

B&RNENS. LONDIN AnGLIC. EdENBURG. MoNSPEL,

ToLos, Fi;.ORENT. Sbc.

SPECIES

M

\$

EXHIBENTES

PLANTAS RITE COGNITAS

AD

GENERA RELATAS,

CUM .

DiFEFERENTIIS SPECEFICISIj

NoMINIBUS TrIVIALIBUS,

Synonymis Selectis,

Locis Natalibus,

Secundum

ST ST EM A SE XVAL B

_,^, DIGESTAS.

>

BOTANICAL,

^ \RDEN. «

TOMUS

f

Editio Scov»qd45 aufta.

Cum Prhikgio S. R M'.tis, Svfcī^ ^ \$, jR, M:tis PtloJii^ ac EhSofit Saxoi^.

HOLMIM

Impensis Direct, LAURENTII SALVII

1763.

[Begin Page: Page 1011]

DIADELPHIA DECANDRIA-

I OII

Cicer fylveare latitolium triphyllum. Bauh.pin, 347.

Cicer fylvellre tertfum. Dod,pempt. 525.

Cicer fvvc/tre verius. L^b. ic x. p.^^.^.

Habitat in Alfthus Helvetici< f, f?-

VexiUufh pubtfiens, AU carina breviores. Calyces €in<

{if braShis 5, cordatiS ^ approximatis^ brcvioribus*

Folia parva^ integsrrii/ia ^ glabra.

* # :^ # •

t^ ONONIS fpicis feffiHhus, legaminibus hutamibus

linearicus pilo.'is. yfj^crw. acd^V). «/r. 42. *.

.Lotus africana annun hirfuta, floribus hiteis. Camm.

hart. 2 />. 163, f. 82.

Hshtat ad Cap. b. rpei. 0.

i§. ONONIS racemis pedunculatis longis , foliis terna capenfis.

m fuborfaiculatjs. Amoen. acad. 6. «/r. 41. *.

Habitat /id Cap, b. fpei. Q.

19. ONONIS pedu.jculis uaifloris mutlcis , fru^ibus cer- redinata-nuts^ ^>his rernatis fubrorundii crenft"s

Anonis pudlla hlrfuta, fiore ex albo & robro color*

mJxto. Morif. hift. z.p. ijij^

Anonis purpurea non fpinofa minor. Barr. u. 2^-4

mala. ' ^^^'

Anonis antloa purtilla, flore purpurafcente. Totiraef.

inli. 408.

^

Hub

ifcofo-pueefceKS , parva , dijfufa , patula. Folfa

\^

petiolata , terrfata ^ orbiculata , cr^mm

crexaia^ vifcaff^'pubffcenfia Stipala ova!es\p^i»U^

-^ ^bt\$tf^ , firrat^. Pedunculi axiiisres ^ umflori^ mutil

ci^ UngUudsne 'vix foliarum. Calyces U>2gitudini

pedunculi. CorollaC aibid^ vexilh f-ubius furpureo^

Calyx defloraius reflectitur apice ad (auUm, Le^U^

)en calyce fesquHongtus , vHlofum. ^

«

ANTHYLLIS,

* Herbace£.

nous hterahcns. Hurt upf. 2.21.* S.ai^: mmfp ..- '*^"P'^>"».

AnthylHs folhs quinato-pianatis^ iasequnribm, floribus

!aterr:hb^{us}. /i*.r/. cliff, g;^ , ^y. /»^m. sgn

Lotus pentaphyllos veficarfa. Bauh, /;». 532.

Trifolfum halicacabum. C^jj. ^j,;^m. lyi t a7

' |