



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789
<https://www.biodiversitylibrary.org/bibliography/4504>

v.1: <https://www.biodiversitylibrary.org/item/23432>

Page(s): Illustration, Title Page, Page 300, Page 301

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 7 January 2025 2:32 AM
<https://www.biodiversitylibrary.org/pdf4/1764225i00023432.pdf>

This page intentionally left blank.

Rare Book
QK 73
G 73
K 48
1789
V. 1
A...
HORTUS KEWENSIS;

OR, A
C A T A L O G U E
OF THE
P L A N T S

CULTIVATED IN THE
ROYAL BOTANIC GARDEN AT KEW.

BY *WILLIAM AITON*,

GARDENER TO

HIS MAJESTY.

IN THREE VOLUMES.]

V O L. I.

MONANDRIA — HEPTANDRIA.

L O N D O N:

PRINTED FOR GEORGE NICOL, BOOKSELLER
TO HIS MAJESTY, PALL MALL.

M. DCC. LXXXIX.

1789

MISSOURI BOTANICAL
GARDEN LIBRARY

Amsonia.

3. T. foliis alternis ovato-lanceolatis, caulibus herbaceis glaberrimis.

Tabernæmontana Amsonia. *Sp. pl.* 308.

Alternate-leav'd Tabernæmontana.

Nat. of North America.

Cult. 1759, by Mr. Philip Miller.

Fl. May and June.

H. 4.

angustifolia.

4. T. foliis linearibus sparsis, caule piloso herbaceo.

Narrow-leav'd Tabernæmontana.

Nat. of North America.

Introd. 1774, by Mr. James Gordon.

Fl. May and June.

H. 4.

C E R O P E G I A. *Gen. pl.* 302.

Contorta. *Folliculi* 2, erecti. *Semina* plumosa. *Corollæ* limbus connivens.

sagittata.

1. C. umbellis subsessilibus, foliis sagittatis. *Linn. mant.* 215.

Arrow-leav'd Ceropegia.

Nat. of the Cape of Good Hope.

Introd. 1775, by Mr. Francis Masson.

Fl.

G. H. h.

D I G Y N I A.

P E R I P L O C A. *Gen. pl.* 303.

Contorta. *Nectarium* ambiens genitalia, filamenta 5 exserens.

græca.

1. P. floribus interne hirsutis terminalibus. *Syst. veget.* 256.

Common

Common Virginian Silk, or Periploca.

Nat. of Syria.

Cult. 1597, by Mr. John Gerard. *Ger. herb.* 754.

Fl. July and August. H. 2.

2. *P. floribus interne hirsutis paniculatis, foliis lanceolato-ellipticis.* *Syst. veget.* 256. *Secamone.*

Green Periploca.

Nat.

Cult. 1775, by John Fothergill, M. D.

Fl. July. G. H. 2.

3. *P. corollis glabris, cymis trichotomis, foliis oblongo-lanceolatis lævibus, caule glabro.* *lævigata.*

Smooth Periploca.

Nat. of the Canary Islands. Mr. Francis Masson.

Introd. 1779.

Fl. G. H. 2.

4. *P. caule hirsuto.* *Sp. pl.* 309. *africana.*

African Periploca.

Nat. of the Cape of Good Hope.

Cult. 1726, by Mr. Philip Miller. *R. S. no.* 209.

Fl. June—September. G. H. 2.

CYNANCHUM. *Gen. pl.* 304.

Contorta. *Nectarium* cylindricum, 5-dentatum.

1. *C. caule volubili perenni aphylo.* *Syst. veget.* 257. *viminale.*

Naked Cynanchum.

Nat. of the Cape of Good Hope.

Cult. 1759, by Mr. Philip Miller. *Mill. dict. edit.* 7.

Euphorbia. 15.

Fl. D. S. 2.

2. C.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Illustration, Title Page]

at ei "See *, fs a ' ' . a
z A . * ae . ;
: ¥ wae ee a. ' ie
- HORTUS KEWE
OR, A
4 so : : ' — Z "
BY WALLIEM ALT OR ¢
GARDENER TO
Pe SEIS MAJESTY, —
ee
I N THR EE vo L
MAJESTY, PALL MALL,
Eeeae, an : en -
Cyt F aie
ae Bos *] *
i
Missour: POTANIGAL
GaR@PN LiBR@RY

[Begin Page: Page 300]

_ 300 PENTANDRIA MONOGYNIA. 'Tabernamontana,
Amfinia. 3. 'T. foliis alternis ovato-lanceolatis, caulibus herbaceis
Ss glaberrimis. ,
Tabernemontana Amfonia. Sp. pi. 308:
Alternate-leav'd Tabernemontana,
Nat. of North America.
Cult-1759, by Mr. Philip Miller. |
Fl, May and June. iY.
angultifo- 4. 'T. foliis linearibus fparfis, caule pilofo herbaceo,
ha, Narrow-leav'd Tabernemontana.
Nat. of North America.
Introd. 1774, by Mr. James Gordon. -
Fl. May and June. HY.
CEROPEGIA. Gen. pl 30% —
Contorta. Folliculi 2, ere&ti. Semina plumofa, G-
| rolle limbis connivens.
fagittata. 1.C. umbellis fubfefeilibus, foliis fagittatis. Linn. matt
ars.
_ ~~ Arrow-leav'd Ceropogia.
Nat. of the Cape of Good Hope.
Introd. 1775, by Mr. Francis Maffon. a
F 7, G. H. he
Dit GY NEE
PERIPLUCA. Gem ph 3d -
Contorta, Neétarium ambiens genitalia, filament
5 exferens. ,
graca. _x. P. floribus interne hifutis terminalibus. Syf vege
256. e

: Common

[Begin Page: Page 301]

PENTANDRIA DIGYNrAa. Periploca.

Common Virginian Silk, or Periploca.

Nat. of Syria.

Cult. 1597, by Mr. John Gersed: Ger. herb. 754:

Fi, July and Augutt. H. kh.

2: P. floribus interne hirfutis paniculatis, foliis lanceolato-
ellipticis, Sy/f. veget. 256.

Green Periploca.

Nat.

Cult. 1775, by John Fothergill, M.D.

Fl. July. G. Heb.

3: P. corollis glabris, cymis trichotomis, foliis oblongo-
lanceolatis lzvibus, caule glabro. »

Smooth Periploca.

Nat. of the Canary Iflands. Mr. Francis Maffon.

Introd. 1779. 7

<= G. Hb.

4. P. caule hirfuto. Sp. pi. 309.

African Periploca.

Nat. of the Cape of Good Hope.

Cult. 1726, by Mr. Philip Miller. R. S. no. 209.

_ #1 June—September. G. H. hs

CYNANC HUM. Gen. pl. 304-

_ Contorta. Neétarium cylindricum, 5-dentatum.

i. c. caule volubili perenni aphylo. Syft. veget. 2576

Naked Cynanchum.

Nat. of the Cape of Good Hope.

Cult. 1759, by Mr. Philip Miller. AGU. did. edit. 7.

Euphorbia. 15,

fi, D, Sis

2s cS

gor

Secamone.

'levigata.

africana.

viminale.