



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789
<https://www.biodiversitylibrary.org/bibliography/4504>

v.1: <https://www.biodiversitylibrary.org/item/23432>

Page(s): Illustration, Title Page, Page 14

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 8 April 2024 2:46 AM
<https://www.biodiversitylibrary.org/pdf4/1688550i00023432.pdf>

This page intentionally left blank.

Rare Book
QK 73
G 73
K 48
1789
V. 1
A...
HORTUS KEWENSIS;

OR, A

C A T A L O G U E

OF THE

P L A N T S

CULTIVATED IN THE

ROYAL BOTANIC GARDEN AT KEW.

BY *WILLIAM AITON*,

GARDENER TO

HIS MAJESTY.

IN THREE VOLUMES.

V O L. I.

MONANDRIA — HEPTANDRIA.

L O N D O N:

PRINTED FOR GEORGE NICOL, BOOKSELLER
TO HIS MAJESTY, PALL MALL.

M. DCC. LXXXIX.

1789

MISSOURI BOTANICAL
GARDEN LIBRARY

americana.

3. *O.* foliis lanceolato-ellipticis integerrimis, racemis angustatis, bracteis omnibus persistentibus connatis parvis.

Olea americana. *Linn. mant.* 24.

American Olive.

Nat. of Carolina and Florida.

Cult. 1758, by Mr. Philip Miller.

Fl. June.

G. H. h.

excelsa.

4. *O.* foliis ellipticis integerrimis, racemis angustatis, bracteis perfoliatis: baseos cyathiformibus persistentibus; superioribus deciduis magnis foliaceis.

Laurel-leav'd Olive.

Nat. of Madeira. Mr. Francis Masson.

Introd. 1784.

Fl.

G. H. h.

fragrans.

5. *O.* foliis lanceolatis serratis, pedunculis lateralibus aggregatis unifloris. *Thunb. japon.* 18. tab. 2. *Syst. veget.* 57.

Quaj-fa. *Osbeck.* 250.

Sweet-scented Olive.

Nat. of CochinChina, China, and Japan.

Introd. 1771, by Benjamin Torin, Esq.

Fl. July and August.

G. H. h.

CHIONANTHUS. *Gen. pl.* 21.

Cor. 4-fida: laciniis longissimis. *Drupæ* nucleus striatus.

virginica.
latifolia.

1. *C.* pedunculis trifidis trifloris. *Sp. pl.* 11.

α foliis ovato-ellipticis.

Broad-leav'd Fringe-tree.

angustifolia.

β foliis lanceolatis.

Narrow-leav'd Fringe-tree.

Nat.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Illustration, Title Page]

at ei "See *, fs a ' ' . a
z A . * ae . ;
: ¥ wae ee a. ' ie
- HORTUS KEWE
OR, A
4 so : : ' — Z "
BY WALLIEM ALT OR ¢
GARDENER TO
Pe SEIS MAJESTY, —
ee
I N THR EE vo L
MAJESTY, PALL MALL,
Eeeaeene, an : en -
Cyt F aie
ae Bos *] *
i
Missour: POTANIGAL
GaR@PN LiBR@RY

[Begin Page: Page 14]

ee,
14 DIANDRIA MONOGYNta. Olea.
america- 3, O. foliis lanceolato-ellipticis integerrimis, racemis an-
Nae gutatis, braCteis ommibus perfifentibus connatis
parvis.
Olea americana, Linn. mant. 24.
American Olive,
Nat, of Carolina and Florida;
Cult. 1758, by Mr. Philip Miller, faa 1 {Eg
Fi, June, ; G.H. bh .
excelfa, . 4.Q. foliis ellipticis intenersimnis: racemis anguttatis,
: braéteis perfoliatis: bafeos cyathiformibus _perfif-
_ | -tentibus; fuperioribus deciduis magnis foliaceis. —
Laurel-leav'd Olive,
Nat. of Madeira. Mr. Francis Ma ott,
eee Sc! eee |
Fi, : aS ee
fragrans. 5. O. foliis lanceolatis ferratis, pedunculis lateralibus ag-
Ss ~ gregatis unifloris,. Thunb. aber fab 2. Syft.
veget. 57.
Quaj-fa. Of). it. 250.
Sweet-fcented Olive. . —
. Nat. of Cochinchina, Chin, sal aa
ae Introd. 1771, by Benjamin Torin, Ef.
Fi. July and Auguft. G.H. b.
. CHIONANTHUS, Gen. pl. 21.
Cor. 4-fida: laciniis longiffimis, Drupe nucleus
. ftriatus, ~

virginica, 1,C, peduticalis trifidis trifloris. Sp. pl 11.

latifolia. — & foliis ovato-elli

Broad-leav'd Fri ringe-tree,

8 foliis-lanceolatis, .

Narhaw-ay 4 F ringe-tree, =

. Nat.

angutti-