



<https://www.biodiversitylibrary.org/>

**Supplementum plantarum Systematis vegetabilium
editionis decimae tertiae, Generum plantarum editionis
sextae, et Specierum plantarum editionis secunda**

Brunsvigae, Impensis Orphanotropei, 1781

<https://www.biodiversitylibrary.org/bibliography/555>

Item: <https://www.biodiversitylibrary.org/item/10321>

Page(s): Title Page, Page 211

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 25 January 2025 3:22 AM

<https://www.biodiversitylibrary.org/pdf4/1769255i00010321.pdf>

This page intentionally left blank.

C 574. (13)

Linnaeus
QK 41
C 574
1781

SUPPLEMENTUM PLANTARUM

SYSTEMATIS VEGETABILIIUM

Editionis decimæ tertice,

GENERUM PLANTARUM

Editionis sextæ,

et

SPECIERUM PLANTARUM

Editionis secundæ.

Editum

A

CAROLO a LINNÉ,

Med. Doct.

Med. & Botan. Prof. Upsal.

Hort. Botan. Præf.

BRUNSVIGÆ,
Impensis Orphanotrophei,

1781.

MISSOURI
BOTANICAL
GARDEN.

tetrapetalus. LORANTHUS floribus ad articulos quatuor verticillatis tetrapetalis, foliis oblongis oppositis subsessilibus.

Habitat in Nova Zeelandia. *Equ. Bäck. h.*

Folia opposita, subpetiolata, oblonga, integerrima.

Flos utrinque ad latera foliorum unus, eoque ad singulum articulum quatuor, & sic per totum caulem.

falcatus. LORANTHUS racemis paucifloris axillaribus, foliis linearibus obtusis lateraliter falcatis glaucis.

Habitat in Madrafs, arboribus parasitica. *König. h.*

Folia margine deorsum vergente falcata, integerrima, avenia, cana.

Racemi axillares, pauciflori,

Flores falcati, fulvi.

DIGYNIA.

repens. FALKIA.

Habitat in Cap. bonæ spei inundatis. *Thunberg.*

Herba & structura floris omnino Convolvuli, sed femina nimis differunt, ut cum Convolvulis combinari queat.

procera. GAHNIA.

Gahnia procera. *Forst. gen. n. 26. Forst. Abt. upf. v. 3. p. 178.*

Habitat in Novæ Zeelandiæ collibus. *Sparmann & Forster.*

Gramen semiorgyale & ultra.

Schoenis africanis simillima, & forte jure illius generis.

TRIGYNIA.

bipinnatus. RUMEX floribus dioicis, foliis bipinnatis.

Habitat in Marocco. *U.*

Caules spithamæi, adscendentes, læves.

Folia cordata, bipinnatifida: laciniis fere ad costam distinctis, singulis pinnatis s. dichotomis.

Stipulæ membranaceæ, diaphanæ, albæ.

Racemas terminalis, compositus: racemulis simplicibus.

Calyx obtusus.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Title Page]

SUPPLEMENTUM
PLANTARUM
SYSTEMATIS VEGETABILIIUM
Editionis decime tertie,
GENERUM PLANTARUM
* . E
ERA us iini.
Editum
A
CAROLO a LINNÉ, `
Med. Dot,
Med, & Botan. Prof. Upfal,
Hort, Botan, Pref.
BRUNSVIG Æ,
]mpenfis Orphanotrophei,
T7323 15

[Begin Page: Page 211]

ker LDRANTHUS fem uie articüle
libus. TEMO Ee
E ulis zm fübpetiolata nga, integerrima.
Flos utrinque ad latera foliorum unus, eoque ad fin-
gu decimam € & fic per totum cau-
cai P LORANTHUS racemis paucifloris axillaribus, fo-
=> => — füis linearibus obtufis lateraliter falcatis glaucis.
Habitat in Madraís, arboribus parafitica. König. h.
Folia margine deorfum vergente falcata , integerrima,
avenia, cana.
Racemi axillares ra
; en ie
FALKIA. ; um
Habitat in Cap. Tehe bei inundatis, Thunberg.
Herba & fiructura floris omnino Convolvuli, fed femi-
`na nimis differunt, ut cum Convalvulis combina-
ri queat, É
procera. GAHNIA.
Gahnia procera. Forf. gen.» n. Aa Forf. T fos
a neris,
e | DIMIYMA d
^u E E ATRA laciniis. D e ad E
tis, fing A E d otomi