



<https://www.biodiversitylibrary.org/>

**Madroño; a West American journal of botany**

Berkeley, California Botanical Society,

<https://www.biodiversitylibrary.org/bibliography/65344>

**v.42:no.1-4 (1995:Jan.-Oct.):**

<https://www.biodiversitylibrary.org/item/185854>

Page(s): Title Page, Page 422

Holding Institution: Smithsonian Libraries and Archives

Sponsored by: Biodiversity Heritage Library

Generated 12 April 2024 2:48 AM

<https://www.biodiversitylibrary.org/pdf4/1690909i00185854.pdf>

This page intentionally left blank.




# MADROÑO

A WEST AMERICAN JOURNAL OF BOTANY

## Contents

- WIND-SHELTERS AS SAFE SITES FOR ESTABLISHMENT OF *LUPINUS ARBOREUS*, A COASTAL SPECIES  
*Barbara L. Gartner* 1
- ERIGERON BISTIENSIS* (ASTERACEAE: ASTEREA): A NEW SPECIES FROM NORTHWESTERN NEW MEXICO  
*Guy L. Nesom and Bill Hevron* 12
- OBSERVATIONS OF *PINUS MAXIMARTINEZII* RZED  
*Jeffrey K. Donahue and Carlos Mar Lopez* 19
- A NEW SPECIES OF *QUERCUS* (FAGACEAE, SECT. *LOBATAE*, GROUP *RACEMIFLORAE*) FROM THE SIERRA MADRE OCCIDENTAL, MEXICO  
*R. Spellenberg, J. R. Bacon, and D. E. Breedlove* 26
- SESSILE-FLOWERED SPECIES IN THE *NAVARRETIA LEUCOCEPHALA* GROUP (POLEMONIACEAE)  
*Alva G. Day* 34
- GAP ANALYSIS OF THE ACTUAL VEGETATION OF CALIFORNIA 1. THE SOUTHWESTERN REGION  
*Frank W. Davis, Peter A. Stine, David M. Stoms, Mark I. Borchert, and Allan D. Hollander* 40
- NOTES
- THE STATUS OF *CLARKIA MOSQUINII* (ONAGRACEAE)  
*L. D. Gottlieb and Lawrence Janeway* 79
- CUSHION-LIKE FRUTICOSE LICHENS AS *DUDLEYA* SEED TRAPS AND NURSERIES IN COASTAL COMMUNITIES  
*Richard E. Riefner, Jr. and Peter A. Bowler* 81
- NOTEWORTHY COLLECTIONS
- ARIZONA 83
- UTAH 83
- WASHINGTON 84
- REVIEWS
- INTERMOUNTAIN FLORA. VASCULAR PLANTS OF THE INTERMOUNTAIN WEST, U.S.A. VOLUME 5. ASTERALES, BY ARTHUR CRONQUIST  
*David J. Keil* 85
- ETHNOBOTANY OF THE CALIFORNIA INDIANS. VOLUMES 1 AND 2, BY B. M. BECK AND S. S. STRIKE  
*Peggy L. Fiedler* 88
- CONSERVATION BIOLOGY: THE THEORY AND PRACTICE OF NATURE CONSERVATION, PRESERVATION, AND MANAGEMENT, BY P. L. FIEDLER AND S. K. JAIN (EDS.)  
*Beth Braker* 89
- ANNOUNCEMENT 18, 39, 91



PUBLISHED QUARTERLY BY THE CALIFORNIA BOTANICAL SOCIETY



Lamiaceae and Verbenaceae includes 1) a table comparing the two families as they are circumscribed in the book, and 2) a branching diagram illustrating the relative positions of groups of Mints and Verbs. This explanatory approach allows the reader to understand the reasoning behind statements of relationship, rather than having to accept the dogma of relationships that exists in many taxonomic texts.

There is a richly illustrated glossary at the end of the book, along with one appendix that covers Cronquist's scheme of family relationships, and a chart summarizing the characteristics of the families covered in the book. The latter will be especially appreciated by students.

The only problem we had with the book is based on our western North American perspective. We are dismayed that the Hydrophyllaceae, Garryaceae, Fouquieriaceae, and Limnanthaceae were not included. We were also disappointed that the "little aquatic monocots" (e.g., Potamogetonaceae, Zanicelliaceae, Zosteraceae, Hydrocharitaceae, Juncaginaceae) were left out—all the more dismaying because Zomlefer's drawings of these families would be a great aid to those of us who are not aquatically-minded. We hereby make a plea here for future editions to include families that are important in the west, even if they aren't well-represented in the east.

A plant taxonomy course where family relationships are a component is sure to benefit from Zomlefer's book. It does more than list characters—it stimulates interest. The book will also serve professional and lay botanists as a reference on family characteristics and relationships. To be sure, as systematic research continues to resolve relationships among families, parts of this book will become dated. Nevertheless, Zomlefer has provided a first step toward a modern text in plant taxonomy. Finally, it is inexpensive, a bargain at less than \$30.

—ROBERT PATTERSON AND THE STUDENTS IN THE 1995 PLANT TAXONOMY CLASS,  
Department of Biology, San Francisco State University, San Francisco, CA 94132.

*Revision of the Genus Sambucus. Dissertation Botanica*, vol. 223, pp. [i-ii], 1-227-  
[256]. By RICHARD BOLLI. 1994. From E. Schweizerbart'sche Verlagsbuchhandlung,  
Johannesstrasse 3A, D-70176 Stuttgart 1, Germany. Softcover US\$45.51. ISBN 3-443-  
64135-0.

This new worldwide monograph (the work is much more than a mere "revision") on *Sambucus* uses morphology, anatomy, ecology, karyology, and biochemistry to recognize nine species (*S. ebulus* with 2 subspp.; *S. wightiana*; *S. adnata*; *S. gaudi-chaudiana*; *S. australasica*; *S. javanica*; *S. nigra* with 6 subspp.; *S. australis*; *S. racemosa* with 2 var.), eight subspecies, and two varieties. Bolli reduces some commonly accepted species to subspecies or varieties. For instance, relevant to *The Jepson Manual's* (1993) treatment of Californian taxa of *Sambucus*, according to Bolli *S. melanocarpa* becomes *S. racemosa* var. *melanocarpa*, *S. mexicana* becomes *S. nigra* subsp. *canadensis*, and *S. racemosa* var. *microbotrys* (this is not in Bolli's synonymy) and var. *racemosa* become *S. racemosa* var. *racemosa*. Bolli's two evolutionary scenarios both postulate "herbaceous *Sambucus* . . . , probably having evolved in Central Asia, . . . to represent the ancestral group" leading "independently" to "the woody taxa of both the Far East and the Western Hemisphere." *Viburnum* and *Adoxa* seem most closely related, but Bolli favors Sambucaceae excluded from Caprifoliaceae and Dipsacales. This important dissertation done at Zürich should not be overlooked due to the rather low circulation of *Dissertationes Botanicae*.

—RUDOLF SCHMID, Department of Integrative Biology, University of California,  
Berkeley, California 94720-3140.



The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Title Page]

VOLUME 42, NUMBER 1

JANUARY-MARCH 1995

MADRONO

A WEST AMERICAN JOURNAL OF BOTANY

Contents

Wind-Shelters as Safe Sites for establishment of *Lupinus arboreus*, a

Coastal Species

Barbara L. Gartner

*Erigeron bistiensis* (Asteraceae: Astereae): A New Species From Northwestern

New Mexico

Guy L. Nesom and Bill Hevron

Observations of *Pinus maximartinezii* Rzed

Jeffrey K. Donahue and Carlos Mar Lopez

A New Species of *Quercus* (Fagaceae, Sect. *Lobatae*, Group *Racemiflorae*) from

THE Sierra Madre Occidental, Mexico

R. Spellenberg, J. R. Bacon, and D. E. Breedlove

Sessile-Flowered Species in the *Navarretia leucocephala* Group (Polemoniaceae)

Alva G. Day

Gap Analysis of the Actual Vegetation of California 1 . The Southwestern

Region

Frank W. Davis, Peter A. Stine, David M. Stoms, Mark I. Borchert, and Allan D. Hollander

NOTES

The Status of *Clarkia mosquinii* (Onagraceae)

L. D. Gottlieb and Lawrence Janeway

Cushion-like Fruticose Lichens As Dudleya Seed Traps and Nurseries in

Communities in the Colorado Plateau

Richard E. Riefner, Jr and Peter A. Bowler

NOTEWORTHY COLLECTIONS / a'

Arizona / \ I

Utah V

Washington Li BR/^^\*^

REVIEWS

Intermountain Flora. Vascular Plants of the Intermountain West, U.S.A.

Volume 5. Asterales, by Arthur Cronquist

David J. Keil

Ethnobotany of the California Indians. Volumes 1 and 2,

BY B. M. Beck and S. S. Strike

Peggy L. Fiedler

Conservation Biology: The Theory and Practice of Nature Conservation,

Preservation, and Management, by P. L. Fiedler and S. K. Jain (eds.)

Beth Braker 89

ANNOUNCEMENT 18,39,91

12

19

26

34

40

79

83

83

84

**[Begin Page: Page 422]**

422

## MADRONO

[Vol. 42]

Lamiaceae and Verbenaceae includes 1) a table comparing the two families as they are circumscribed in the book, and 2) a branching diagram illustrating the relative positions of groups of Mints and Verbs. This explanatory approach allows the reader to understand the reasoning behind statements of relationship, rather than having to accept the dogma of relationships that exists in many taxonomic texts.

There is a richly illustrated glossary at the end of the book, along with one appendix that covers Cronquist's scheme of family relationships, and a chart summarizing the characteristics of the families covered in the book. The latter will be especially appreciated by students.

The only problem we had with the book is based on our western North American perspective. We are dismayed that the Hydrophyllaceae, Garryaceae, Fouquieriaceae, and Limnanthaceae were not included. We were also disappointed that the "little aquatic monocots" (e.g., Potamogetonaceae, Zanichelliaceae, Zosteraceae, Hydrocharitaceae, Juncaginaceae) were left out— all the more dismaying because Zomlefer's drawings of these families would be a great aid to those of us who are not aquatically-minded. We hereby make a plea here for future editions to include families that are important in the west, even if they aren't well-represented in the east.

A plant taxonomy course where family relationships are a component is sure to benefit from Zomlefer's book. It does more than list characters— it stimulates interest. The book will also serve professional and lay botanists as a reference on family characteristics and relationships. To be sure, as systematic research continues to resolve relationships among families, parts of this book will become dated. Nevertheless, Zomlefer has provided a first step toward a modern text in plant taxonomy. Finally, it is inexpensive, a bargain at less than \$30.

—Robert Patterson and the Students in the 1995 Plant Taxonomy Class,  
Department of Biology, San Francisco State University, San Francisco, CA 94132.  
Revision of the Genus *Sambucus*. Dissertation *Botanicce*, vol. 223, pp. [i-ii], 1-227-  
[256]. By Richard Bolli. 1994. From E. Schweizerbart'sche Verlagsbuchhandlung,  
Johannesstrasse 3A, D-70176 Stuttgart 1, Germany. Softcover US\$45.51. ISBN 3-443-  
64135-0.

This new worldwide monograph (the work is much more than a mere "revision") on *Sambucus* uses morphology, anatomy, ecology, karyology, and biochemistry to recognize nine species (*S. ebulus* with 2 subspp.; *S. wightiana*; *S. adnata*; *S. gaudichaudiana*; *S. australasica*; *S. javanica*; *S. nigra* with 6 subspp.; *S. australis* \ *S. racemosa* with 2 var.), eight subspecies, and two varieties. Bolli reduces some commonly accepted species to subspecies or varieties. For instance, relevant to The Jepson Manual (1993) treatment of Californian taxa of *Sambucus*, according to Bolli *S. melanocarpa* becomes *S. racemosa* var. *melanocarpa*, *S. mexicana* becomes *S. nigra* subsp. *canadensis*, and *S. racemosa* var. *microbotrys* (this is not in Bolli's synonymy) and var. *racemosa* become *S. racemosa* var. *racemosa*. Bolli's two evolutionary scenarios both postulate "herbaceous *Sambucus* . . . , probably having evolved in Central Asia, ... to represent the ancestral group" leading "independently" to "the woody taxa of both the Far East and the Western Hemisphere." *Viburnum* and *Adoxa* seem most closely related, but Bolli favors *Sambucaceae* excluded from *Caprifoliaceae* and *Dipsacales*. This important dissertation done at Zurich should not be overlooked due to the rather low circulation of *Dissertationes Botanicce*.

—Rudolf Schmid, Department of Integrative Biology, University of California,  
Berkeley, California 94720-3140.