



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated  
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789  
<https://www.biodiversitylibrary.org/bibliography/4504>

**v.2:** <https://www.biodiversitylibrary.org/item/23433>

Page(s): Illustration, Title Page, Page 341, Page 342

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 12 April 2024 9:34 AM  
<https://www.biodiversitylibrary.org/pdf4/1691006i00023433.pdf>

This page intentionally left blank.

*Rare Books*  
A  
Q K73  
G73  
K48  
1789  
V.2

*HORTUS KEWENSIS;*

O R, A

C A T A L O G U E  
O F T H E  
P L A N T S

CULTIVATED IN THE  
ROYAL BOTANIC GARDEN AT KEW,

B Y W I L L I A M A I T O N,  
GARDENER TO  
H I S M A J E S T Y.

I N T H R E E V O L U M E S.]

---

V O L. II.

OCTANDRIA — MONADELPHIA.

---

L O N D O N:

PRINTED FOR GEORGE NICOL, BOOKSELLER  
TO HIS MAJESTY, PALL MALL.

M,DCC,LXXXIX.

1789  
MISSOURI BOTANICAL  
GARDEN LIBRARY

DIDYNAMIA ANGIOSPERMIA. Scrophularia. 341

3. S. foliis cordatis petiolatis decurrentibus obtusis, caule membranis angulato, racemis terminalibus. Sp. pl. 864. *Curtis lond.* *aquatica.*  
 Water Figwort.  
*Nat.* of Britain.  
*Fl.* May—July. H. 4.
4. S. foliis cordatis subtus tomentosis basi appendiculatis, racemis terminalibus. Syst. veget. 560. *auriculata.*  
 Ear'd-leav'd Figwort.  
*Nat.* of Spain.  
*Introd.* 1772, by Mons. Richard.  
*Fl.* July. H. 4.
5. S. foliis cordatis duplicato-ferratis pubescentibus, paniculis terminalibus trichotomis foliis interstinctis. *Scorodonia.*  
 Scrophularia Scorodonia. Sp. pl. 864.  
 Balm-leav'd Figwort.  
*Nat.* of England.  
*Fl.* May—July. H. 4.
6. S. foliis oblongo-lanceolatis cordatis duplicato-ferratis glabratæ, glabris, paniculis racemosis terminalibus trichotomis, caule suffruticoso.  
 Spear-leav'd Figwort.  
*Nat.* of the Canary Islands, Mr. Francis Masson.  
*Introd.* 1779.  
*Fl.* April and May. G. H. ♂.
7. S. foliis cordatis oblongis dentatis: dentibus integerimis: baseos profundioribus. Syst. veget. 560. *betonicifolia.*  
 Betony-leav'd Figwort.  
*Nat.* of Spain.  
*Cult.* 1739, by Mr. Ph. Miller, Mill. diet, vol. 2. n. 3.  
*Fl.* June—August. H. 4.

*orientalis.* 8. S. foliis lanceolatis ferratis petiolatis: caulinis ternis; rameis oppositis. *Sp. pl.* 864.

Hemp-leav'd Figwort.

*Nat.* of the Levant.

*Cult.* 1712. *Philosoph. transact.* n. 333. p. 416.  
n. 60.

*Fl.* July and August.

H, 4.

*frutes-*  
*cens,*

9. S. foliis subcarnosis sessilibus levibus apice recurvis,  
*Syst. veget.* 560.

Shrubby Figwort.

*Nat.* of Portugal,

*Cult.* 1768, by Mr. Ph. Miller. *Mill. diet. edit.* 8,

*Fl.* June—August.

G. H. h,

*vernalis.*

10. S. foliis cordatis pubescentibus duplicato-serratis,  
paniculis axillaribus dichotomis.

Scrophularia vernalis. *Sp. pl.* 864.

Yellow Figwort.

*Nat.* of England.

*Fl.* March—May.

H, 3.

*arguta.*

11. S. foliis cordatis glabris duplicato-serratis, paniculis  
axillaribus dichotomis, capsulis acuminatis.

Slender upright Figwort.

*Nat.* of Madeira and Teneriffe. Mr. Fr. Masson.

*Introd.* 1778.

*Fl.* May and June.

G. H. o.

OBS. Differt a Scr. vernali caule foliisque glabris, et  
floribus minoribus rubris.

*sambuci-*  
*folia.*

12. S. foliis interrupte pinnatis cordatis inæqualibus, ra-  
cemo terminali: pedunculis axillaribus geminis  
dichotomis. *Sp. pl.* 865.

Elder-leav'd Figwort.

*Nat.*

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Illustration, Title Page]

HORTUS KEWENSIS;  
OR, A ae  
-€ £T ASO G UE  
Or THE |  
PE ji A nN re  
CuiniraTks th-Fas  
ROYAL BOTANIC GARDEN AT KEW,  
BY WIELELLAM ALLO  
GARDENER TO  
iis MATE SCY,  
¥- 0. IL, 2  
OCTANDRIA— MONADELPHIA.  
LONDON:  
PRINTED FOR GEORGE NICOL, SOOKSRERER  
TO HIS MAJESTY, PALL MALL.  
M,DCC,LXXXIX, © ;  
it Ch = eee  
" Missour: BoPaNICAL. ———  
\_ GARDEN LIBRARY : .

[Begin Page: Page 341]

DIDYNAMIA ANGIOSPERMIA.- Scrophularia. 342  
3. S. foliis cordatis petiolatis decurrentibus obtusis, caule aquatica,  
membranis angulato, racemis terminalibus. Sp. p/.  
864. Curtis lond.  
Water Figwort.  
Nat. of Britain,  
Fi, May July. Soap  
4. S. foliis cordatis fubtus tomentofis bafiappendiculatis, aurigula+  
racemis terminalibus. Sy/t.- veget. 560. L\* ta.  
Far'd-leav'd Figwort. .  
Nat. of Spain.  
Introd. 1772, by Monf. Richard,  
Fi, July. H, 4,  
5. S. foliis cordatis duplicato-ferratis pubefcentibus, pa- Scorado=.  
niculis terminalibus trichotomis foliis interftin@tis, #4  
Scrophularia Scorodonia. Sp. pl. 804.  
Balm-leav'd Figwort,  
Naz. of England. :  
Fl. May—July. H. 4.  
§. S. foliis oblongo-lanceolatis cordatis duplicato-ferratis : glabrata,  
glabris, paniculis racemofis terminalibus trichoto= %  
mis, caule fuffrutofo.  
Spear-leav'd Figwort.  
Nat. of the Canary Iflands, Mr. Francis Maffon.  
Introd, 1779.  
Fl, April and May, 'G.H, é.  
7: S. foliis cordatis oblongis dentatis: dentibus integer-, betonici-

rimis: bafeos profundioribus. Sy/?, veget. 560, fi 'lia.  
Betony-leav'd Fi igwort.  
Nat. of Spain,  
Cult. 1739, by Mr. Ph. Miller, Mill. di€t, val. 2« #- 3-  
Fi, June—Augutt, HL.  
23 oe 3 28:5,

[Begin Page: Page 342]

342 DIDYNAMIA ANGIOSPERMIA, Scrophularia,  
134 0 ¢@ 6  
orientalis, 8. S, foliis Yanceolatis ferratis. petiolatis caulinis tere i  
weyers nis \$" rameis — Sp. pl 864, see 9 i eat  
\_Hemp-leav'd Figwort, ae  
Nat, of the Vevaas:  
Cult. 1712. Philofoph, 'nefits n 333 p. 6  
n. 60,  
Fi. July and Auguft. Hy ys  
Srutef- 9g. 5. foliis fubcarnofis feffilibus levibus apice recurviss oi  
fensy Sift, vegtt, \$60, : |  
Shrubby Figwort.  
Nat. of Portugal, As  
Cult, 1768, by Mr. Ph. Miller, 4ll, didi, edit. a Hes  
Y FL June—Auguft. G. Hy isp  
wernalis, 10, 8, foliis cordatis pubefcentibus Suey ,  
sic paniculis axillaribus dichotomis.  
Scrophularia vernalis, Sp. ss 864, a a  
Yellow Figwort, ee  
Nat. of England, vr ae  
« Fl. March—May.  
ss arguta, © 41,8. foliis cordatis glabris duplicito-ferratis, | pen wn  
ae \_ ~~ axillaribus dichotomis, capfulis acumiriatisy = ee =  
Slender Moke Figwort, —  
— fers minoribus rubris,:  
32, 6. foliis interrupte pinnatis cordatis ineequ lib  
cemo terminali: pedunculis axil Haribus  
dichotomis. Sp. pi, 865. oe  
| Elder-leav'd Figwort. —  
be FP .gianoaiaale  
epee ese grins  
§ sf '  
| : fambuci~