



<https://www.biodiversitylibrary.org/>

**Supplementum plantarum Systematis vegetabilium
editionis decimae tertiae, Generum plantarum editionis
sextae, et Specierum plantarum editionis secunda**

Brunsvigae, Impensis Orphanotropei, 1781

<https://www.biodiversitylibrary.org/bibliography/555>

Item: <https://www.biodiversitylibrary.org/item/10321>

Page(s): Title Page, Page 36, Page 37, Page 251

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

Generated 23 April 2024 6:08 AM

<https://www.biodiversitylibrary.org/pdf4/1693939i00010321.pdf>

This page intentionally left blank.

C 574. (13)

Linnaeus
QK 41
C 574
1781

SUPPLEMENTUM PLANTARUM

SYSTEMATIS VEGETABILIIUM

Editionis decimæ tertice,

GENERUM PLANTARUM

Editionis sextæ,

et

SPECIERUM PLANTARUM

Editionis secundæ.

Editum

A

CAROLO a LINNÉ,

Med. Doct.

Med. & Botan. Prof. Upsal.

Hort. Botan. Præf.

BRUNSVIGÆ,
Impensis Orphanotrophei,

1781.

MISSOURI
BOTANICAL
GARDEN.

DODECANDRIA.

MONOGYNIA.

1392. DODECAS.

*Charakter essentialis.**Calyx* semiquadrifidus, corollifer, superus. *Corolla* pentapetala. *Capsula* unilocularis, calyce connata.*Charakter naturalis.***CAL.** *Perianthium* superum, monophyllum, turbinatum, semiquadrifidum.*Laciniae* ovatae, patentes.**COR.** *Petala* quatuor, sessilia, calyci inserta, subrotunda.**STAM.** *Filamenta* duodecim, capillaria, calyce breviora, receptaculo inserta.*Antherae* oblongae.**PIST.** *Germen* semifuperum.*Stylus* filiformis, staminibus longior.*Stigma* simplex.**PER.** inferum, ovatum, uniloculare, calyce patulo coronatum, intra quem apex capsulae denudatus quadri-valvis.**SEM.** numerosa, oblonga minuta.*Obs.* Forte *Jussieuæ* species.

TRIGYNIA.

1393. VISNEA.

*Charakter essentialis.**Corolla* pentapetala. *Calyx* pentaphyllus, inferus. *Stigmata* tria.*Nux* bi- s, trilocularis, semi infera.*Charakter naturalis.***CAL.** *Perianthium* inferum, pentaphyllum, persistens.*Foliola* lanceolata, recurvata: tria exteriora pilosa.**COR.** pentapetala.*Petala* elliptica, aequalia, patentia, calyce vix longiora, integra.**STAM.** *Filamenta* duodecim, erecta, receptaculo inserta, filiformia, petalis breviora.*Antherae* quadrangulae, erectae, arista terminatae.**PIST.**

PIST. *Germen* superum, hirtum, attenuatum quasi in Sty-
lum hirtum brevissimum.

Styli tres, filiformes, glabri.

Stigmata simplicia.

PER. *Nux* ovata, glabra, acuminata, bi- s. trilocularis, se-
miinfera, calycis foliolis conniventibus inclusa,
& per maximam partem monophylla calycis par-
te tecta, quæ cum nucis connata, at superior
tertia pars nucis intra calycem nuda.

SEM. singulo loculamento unicum.

*In memoriam Dni Visne, Portugalli, amore & cognitione
plantarum noti.*

1394. TACCA. *Forst. gen. n. 35.*

Charakter essentialis.

Flos superus. *Corolla* hexapetala, fornicata. *Calyx* he-
xaphyllus. *Bacca* sicca, angulata, trilocularis.

Charakter naturalis.

CAL. *Perianthium* superum, hexaphyllum.

Foliola ovata, conniventia, persistencia.

COR. *Petala* sex, fornicata, æqualia.

STAM. *Filamenta* vix ulla.

Antheræ oblongæ, binæ singulorum petalorum ca-
vo innatæ.

PIST. *Germen* inferum,

Styli tres, breves.

Stigmata obcordata.

PER. *Bacca* sicca, ovata, angulata, trilocularis.

SEM. plura, ovata, striata.

1395. PALLASIA. *Pterococcus. Pallas.*

Charakter essentialis.

Corolla pentapetala. *Calyx* nullus. *Nux* quadrangula-
ris: *angulis* membranaceis, bifidis.

Charakter naturalis.

CAL. nullus.

COR. *Petala* quinque, subrotunda, subinæqualia, persi-
stencia.

DIGYNIA.

decumbens.

AGRIMONIA foliis pinnatis hirsutis, caule procumbente, fructibus undique hispido-uncinatis.

Habitat ad Cap. bonæ spei. 7.

Folia radicalia Potentillæ Anserinæ, pinnata.

Foliola circiter viginti, æqualia, ferrata, ovata, villosissima.

Caules pedales, procumbentes, subaphylli, simplicissimi, transeuntes in racemum.

Fructus ovati, tomentosi, undique adhærentes, stimulis uncinatis, inferi plerumque monospermi, monogyni.

TRIGYNIA.

Mocanera. VISNEA.

Mocanera. *Canar.*

Habitat in sylvis montosis insularum Canariensium.

Franc. Masson. 7.

Frutex parvus.

Caulis subverrucosus, teres.

Folia alterna, erecta, breve petiolata, elliptica, glaberrima, venosa, ferrata, consistentia Lauri.

Pedunculi axillares, solitarii, nutantes, vix petiolo longiores, nudi, uniflori.

Flores parvi.

Corolla lutea.

Flore foecundato erigitur pedunculus, calyx clauditur & incrassatur, eiusque foliola tria exteriora fusca pilosa evadunt.

pinnatifida. TACCA.

Tacca pinnatifida. *Forst. gen. n. 35.*

Tacca littorea. *Rumph. amb. v. 5. p. 328. t. 114.*

Tacca fativa. *Rumph. amb. v. 5. p. 324. t. 112.*

Habitat in India orientali & in Taheiti. *Sparmann.* 7.

Foliis similis Dracontio polyphylo.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Title Page]

SUPPLEMENTUM
PLANTARUM
SYSTEMATIS VEGETABILIIUM
Editionis decime tertie,
GENERUM PLANTARUM
* . E
ERA us iini.
Editum
A
CAROLO a LINNÉ, `
Med. Dot,
Med, & Botan. Prof. Upfal,
Hort, Botan, Pref.
BRUNSVIG Æ,
]mpenfis Orphanotrophei,
T7323 15

[Begin Page: Page 36]

ae DODECANDRIA. Monogynia,
^A DODECANDRIA.
Enn MONOGYNIA. E
H3 DODECAS,
CharaBer Mer.
Is, cerollifer, fūperus. - Corolla pen- ,
iesu, , inci d connata,
sm Pes quatuor, elitin € ze inferas Tn.
` Sram. Filamenta duodecim , capillaria, calyee:breviora, re-
ceptaculo inferta.
. Anthere oblongee, x — RR
: Pist. Germen femifaperum. roe
TUN ER Ais opinor: gior. +
a cand
*
Yu.
Ea dem. Ms
LI
GIAIOYNIAC
1393. VISN EA.
Charatter. effentialis. -
Corolla m a pA be penapiyls, ines, Stig-
Ca
: s Sra ds ua Eb. receptaculo inferta,
" — filiformia, petalis breviora. i\$
; T: ds bes — Y ereCtz, arifta
^

[Begin Page: Page 37]

E -

quafi in Sty-

DODECANDRTA. rigyni:

Pist, Germen faperum, hirtum, attenuati 1

p seg brevi d ux

tyli tres, filif edm ib glabri gv 3 2 =

S ita limplicia. - "SU mot

tigmata

Psr, Nux ovata, glabra, acuminata, wt trilocularis, fe-

aperi lee calycis foliolis rede Mer.

per maximam partem monophy: ycis par-

te te&a, quæ cum nuce connata, at fuperior

^^ tertia pars nucis intra calycem nuda,

se Sula loculamento unicum,

— In memoriam. Dni Vifne, Portugalli, amore €? cojtitione

1 nh nati.

F5 e £f.

| CharaBer PES a F E |

Flos fuperus. Corolla hexapetala, . foina, Chah fe-

xaphyllus. Bacca ficca, angulata; trilocularis.

Charafer miroir, " !

CAL. Perigntà ium fuperum, hexaphyllum.

Ep — - Foliola ovata, Mauri perfiftentia.

Cox. "Petala fex, fornicata, aequalia.

Sram.. | vix ulla. E

2o c Anthera biet, bine 22 Shan petiorum, ca-

— vo innate. - a.

Prst. Germen inferum, 9.

is 2 SH tres, breves © 08 pe 4

.. Stigmata obcordata. (C rta dp

5 PeR. Bacca ficca, ovata, angulata, trilocularis.

[Begin Page: Page 251]

oro

pat

d Mocanera,

pos imm

M^ cxD————

" rome

aor: LS.

Habitat in- [ylvis. pe Dem uis Crarinfm,

Franc. Maffon. À k ,

Frutex parvus,

u

Caulis fabverrucofus, deis

een alterna, erecta, breve petiolata, elli

berrima, xenoix: ferratà, confiftentia donent Lauri; S

(Pedunculi axillares, folitarii , — vix pics

dengue c ee He

-& increffatur, eiusque foliola tra ç exteriora fala

pilofa duni:

TACCA |

~ For, x B. zz

acca soea fio. EZ fs opip v. 5. p: e t lu.

acca fativa, vd amb. v. 5. p. 324. t. TI

labi .in India orientali & in Taheiti' : i

mpa i